

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

19



Europäisches Patentamt

European Patent Office

Office européen des brevets

11 Publication number:

0 145 434
A2

12

EUROPEAN PATENT APPLICATION

21 Application number: 84308386.6

22 Date of filing: 03.12.84

51 Int. Cl.⁴: **C 07 D 211/90**, 07 D 401/12,
C 07 D 409/12, C 07 D 405/12,
A 61 K 31/44
// C07D309/12, C07C69/00,
C07C149/00, C07C91/00,
C07D295/14

30 Priority: 02.12.83 JP 228880/83

43 Date of publication of application: 19.06.85
Bulletin 85/25

64 Designated Contracting States: CH DE FR GB IT LI NL
SE


71 Applicant: OTSUKA PHARMACEUTICAL CO., LTD., 9,
Kandatsukasacho 2-chome, Chiyoda-ku Tokyo 101 (JP)

72 Inventor: Tamada, Shigeharu, 1-16, Aza Nibu Shinkirai
Kitajimacho, Itano-gun Tokushima-ken (JP)
Inventor: Nagami, Kazuyoshi, 21-3, Aza Todoroki
Yoshinari, Ojincho Tokushima-shi (JP)
Inventor: Teramoto, Shuji, 483-10, Kagasuno
Kawauchicho, Tokushima-shi (JP)
Inventor: Tanaka, Tatsuyoshi, 21,
Kitajurokujonishi-3-chome Kita-ku, Sapporo-shi (JP)
Inventor: Nakagawa, Kazuyuki, 774-1, Oomatsu
Kawauchicho, Tokushima-shi (JP)

74 Representative: Ruffles, Graham Keith et al, MARKS &
CLERK 57-60 Lincoln's Inn Fields, London WC2A 3LS
(GB)

54 Novel dihydropyridine derivatives and process for preparing the same.

57 Dihydropyridine derivatives and salts thereof represented
by the general formula,



which possess excellent calcium antagonist effect, hypoten-
sive effect, platelets aggregation inhibitory effect, phospho-
diesterase inhibitory effect, calmodulin inhibitory effect and
peroxidized lipid lowering effect, and thus dihydropyridine de-
rivatives and salts thereof are useful as a coronary blood flow
improving agent such as coronary vasodilator, hypotensive
agent, prophylaxis and treating agents for thrombosis, phos-
phodiesterase inhibitory agent, peroxidized lipid metabolism
lowering agent, anti-inflammatory agent and others.

EP 0 145 434 A2


799P49382

NOVEL DIHYDROPYRIDINE DERIVATIVES
AND PROCESS FOR PREPARING THE SAME

1 FIELD OF THE INVENTION

The present invention relates to dihydro-
pyridine derivatives and salts thereof. More parti-
cularly, the present invention relates to dihydropyridine
5 derivatives and salts thereof, process for preparing the
same, as well as pharmaceutical composition containing
the same as the active ingredient.

The dihydropyridine derivatives and salts
thereof are novel compounds, and are not known in any
10 of prior art references. The dihydropyridine derivatives
and salts thereof according to the present invention are
represented by the general formula (1),



wherein R^1 and R^4 are each a lower alkyl group; R^2 is a
lower alkyl group or a group of the formula $-\text{CH}_2-\text{A}-\text{R}^6$
15 [wherein A is a straight-chain or branched-chain un-
saturated group hydrocarbon group which may have an
oxygen atom or a group of the formula $-\text{N}-\text{R}^7$ (wherein R^7
is a lower alkyl group); and R^6 is a phenyl group which
may have a hydroxyl group]; R^3 is a phenyl group which

- 1 may have 1 to 2 substituents selected from the group consisting of a nitro group, a lower alkyl group which may have 1 to 3 halogen atoms, a lower alkoxy group and a halogen atom; and R^5 is a lower alkyl group, a 1,2,3,6-tetrahydropyridyl-lower alkyl group which may have, as the substituent, a phenyl group which may have halogen atoms or lower alkyl groups as the substituents on the phenyl ring, or a group of the formula $-CH_2-A'-R^8$ [wherein A' is a straight-chain or branched-chain unsaturated hydrocarbon group which may have or may not have an oxygen atom, a sulfur atom, a group of the formula $-N-R^7$ (wherein R^7 is a lower alkyl group) or a group of the formula $-N \begin{array}{c} \diagup \quad \diagdown \\ \text{ } \end{array} N-$ in the unsaturated hydrocarbon group; and R^8 is a phenyl group which may have 1 to 3 substituents selected from the group consisting of a lower alkoxy group, a halogen atom, a lower alkylthio group, a hydroxyl group, a lower alkanoyloxy group, a tetrahydropyranyloxy group and a lower alkoxy-lower alkoxy group; a pyridyl group; a thienyl group; furyl group, or a tetrazolyl group which may have a lower alkyl group as the substituent]; provided that when R^5 is a lower alkyl group, then R^2 should be a group of the formula $-CH_2-A-R^6$ (wherein A and R^6 are the same as defined above).

Dihydropyridine derivatives and salts thereof represented by the general formula (1) possess excellent calcium antagonist effect, hypotensive effect, platelets aggregation inhibitory effect, phosphodiesterase inhibitory effect, calmodulin inhibitory effect and

1 peroxidized lipid lowering effect, and thus the dihydro-
pyridine derivative and salt thereof represented by the
general formula (1) are useful as a coronary blood flow
improving agent such as coronary vasodilator, hypotensive
5 agent, prophylaxis and treating agents for thrombosis,
phosphodiesterase inhibitory agent, peroxidized lipid
metabolism lowering agent, anti-inflammatory agent and
others.

DESCRIPTION OF THE PRIOR ART

10 Compounds similar to the dihydropyridine
derivatives according to the present invention are known
from the disclosures in Japanese Patent Application Kokai
(Laid-open) No. 51-108075 (1976) and Japanese Patent
Application Kokai (Laid-open) No. 56-36455 (1981). Com-
15 pounds of these prior art references are known as useful
hypotensive agents, peripheral and cerebral vasodilating
agents and treating agent for coronary blood vessels.
On the contrary, dihydropyridine derivatives according
to the present invention have features in that they
20 perform their pharmacological effects for longer period
of time with less side-effects as compared with known
compounds. Furthermore, dihydropyridine derivatives
according to the present invention are useful as carcino-
static agents.

25 SUMMARY OF THE INVENTION

An object of the present invention is to provide

1 novel dihydropyridine derivatives.

Another object of the present invention is to provide process for preparing said dihydropyridine derivatives.

5 Further object of the present invention is to provide a pharmaceutical composition containing said dihydropyridine derivative as the active ingredient.

These and other objects and features of the present invention will become more fully apparent from
10 the following description.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Dihydropyridine derivatives represented by the general formula (1) according to the present invention, examples of various substituents as defined in R^1 , R^2 ,
15 R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , A and A' are as follows.

As to the lower alkyl group, an alkyl group having 1 to 6 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, pentyl, hexyl groups and others can be exemplified.

20 As to the lower alkyl group which may have halogen atoms as the substituents, an alkyl group having 1 to 6 carbon atoms which may have 1 to 3 halogen atoms as the substituents, in addition to the above-mentioned alkyl groups having 1 to 6 carbon atoms, trifluoromethyl,
25 2,2-difluoroethyl, 1,1-dichloroethyl, trichloromethyl, dichloromethyl, tribromomethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, 2-chloroethyl, 1,2-dichloroethyl,

1 3,3,3-trichloropropyl, 3-fluoropropyl, 4-chlorobutyl, 3-chloro-2-methylethyl groups and others can be exemplified.

As to the lower alkoxy group, an alkoxy group having 1 to 6 carbon atoms such as methoxy, ethoxy,
5 propoxy, isopropoxy, butoxy, tert-butoxy, pentyloxy, hexyloxy groups and others can be exemplified.

As to the halogen atom, fluorine atom, chlorine atom, bromine atom and iodine atom can be exemplified.

As to the lower alkoxy-lowered alkoxy group,
10 an alkoxyalkoxy group of which the alkoxy moieties having 1 to 6 carbon atoms such as methoxymethoxy, 2-methoxyethoxy, 1-methoxyethoxy, 3-methoxypropoxy, 4-methoxybutoxy, 1,1-dimethyl-2-methoxyethoxy, 5-methoxypentyloxy, 6-methoxyhexyloxy, 2-methyl-3-methoxypropoxy, ethoxymethoxy,
15 3-ethoxypropoxy, 6-ethoxyhexyloxy, 2-propoxyethoxy, 4-propoxybutoxy, 5-butoxypentyloxy, pentyloxymethoxy, 1-pentyloxyethoxy, 1,1-dimethyl-2-hexyloxyethoxy, 3-hexyloxypropoxy groups and others can be exemplified.

As to the lower alkylthio group, an alkylthio
20 group having 1 to 6 carbon atoms, such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, tert-butylthio, pentylthio, hexylthio groups and others can be exemplified.

As to the lower alkanoyloxy group, an alkanoyl-
25 oxy group having 1 to 6 carbon atoms, such as formyloxy, acetyloxy, propionyloxy, butyryloxy isobutyryloxy, penta-noyloxy, tert- butyryloxy, hexanoyloxy groups and others can be exemplified.

1 As to the phenyl group which may have 1 to 3
substituents selected from the group consisting of a
lower alkoxy group, a halogen atom, a lower alkylthio
group, a hydroxyl group, a lower alkanoyloxy group, a tetra-
5 hydropyranyloxy group and a lower alkoxy-lower alkoxy
group, a phenyl group which may have 1 to 3 substituents
selected from the group consisting of an alkoxy group
having 1 to 6 carbon atoms, a halogen atom, an alkylthio
group having 1 to 6 carbon atoms, a hydroxyl group, an
10 alkanoyloxy group having 1 to 6 carbon atoms, a tetra-
hydropyranyloxy group and an alkoxyalkoxy group of which
alkoxy moieties having 1 to 6 carbon atoms, such as
phenyl, 2-, 3- or 4-chlorophenyl, 2-, 3- or 4-fluoro-
phenyl, 2-, 3- or 4-bromophenyl, 2-, 3- or 4-iodophenyl,
15 3,5-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl,
3,4-difluorophenyl, 3,5-dibromophenyl, 2-, 3- or 4-methyl-
thiophenyl, 2-, 3- or 4-ethylthiophenyl, 4-propylthio-
phenyl, 3-isopropylthiophenyl, 2-butylthiophenyl, 4-
hexylthiophenyl, 3-pentylthiophenyl, 4-tert-butylthio-
20 phenyl, 3,4-dimethylthiophenyl, 2,5-dimethylthiophenyl,
2-, 3- or 4-methoxyphenyl, 2-, 3- or 4-ethoxyphenyl, 3-
propoxyphenyl, 4-isopropoxyphenyl, 3-butoxyphenyl, 2-
pentyloxyphenyl, 4-tert-butoxyphenyl, 4-hexyloxyphenyl,
3,4-dimethoxyphenyl, 3,4-diethoxyphenyl, 2,5-dimethoxy-
25 phenyl, 2-, 3- or 4-(2-tetrahydropyranyloxy)phenyl, 2,4-
bis(2-tetrahydropyranyloxy)phenyl, 3-methylthio-4-
chlorophenyl, 2-chloro-6-methylthiophenyl, 2-methoxy-3-

- 1 hydroxyphenyl, 3,4,5-trimethoxyphenyl, 3,4,5-trimethyl-
thiophenyl, 3,4,5-trichlorophenyl, 2-, 3- or 4-hydroxy-
phenyl, 3,4-dihydroxyphenyl, 2,6-dihydroxyphenyl, 3,4,5-
trihydroxyphenyl, 2-, 3- or 4-acetyloxyphenyl, 4-propio-
5 nyloxyphenyl, 3-isopropionyloxyphenyl, 2-butyryloxy-
phenyl, 4-hexanoyloxyphenyl, 3-pentanoyloxyphenyl,
4-tert-butyryloxyphenyl, 3,4-diacetyloxyphenyl, 2,5-
diacetyloxyphenyl, 3,4,5-tridiacetyloxyphenyl, 2-methoxy-
methoxyphenyl, 3-(2-methoxyethoxy)phenyl, 4-(1-methoxy-
10 ethoxy)phenyl, 2-(3-methoxypropoxy)phenyl, 3-(4-
methoxybutoxy)phenyl, 4-(1,1-dimethyl-2-methoxyethoxy)-
phenyl, 2-(5-methoxypentyloxy)phenyl, 3-(6-methoxy-
hexyloxy)phenyl, 4-(2-methyl-3-methoxypropoxy)phenyl,
2-(ethoxymethoxy)phenyl, 3-(3-ethoxypropoxy)phenyl,
15 4-(6-ethoxyhexyloxy)phenyl, 2-(2-propoxyethoxy)phenyl,
3-(4-propoxybutoxy)phenyl, 4-(5-butoxypentyloxy)phenyl,
2-(pentyloxymethoxy)phenyl, 3-(1-pentyloxyethoxy)phenyl,
4-(1,1-dimethyl-2-hexyloxyethoxy)phenyl, 2-(3-hexyloxy-
propoxy)phenyl, 2-(3-hexyloxypropoxy)phenyl groups and
20 others can be exemplified.

As to the phenyl group which may have 1 to 2
substituents selected from the group consisting of a
nitro group, a lower alkyl group which may have 1 to 3
halogen atoms, a lower alkoxy group and a halogen atom,
25 a phenyl group which may have 1 to 2 substituents, on
the phenyl ring, selected from the group consisting of
a nitro group, an alkyl group having 1 to 6 carbon atoms

- 1 which may have 1 to 3 halogen atoms, an alkoxy group
having 1 to 6 carbon atoms, and a halogen atom, such as
a phenyl, 2-, 3- or 4-chlorophenyl, 2-, 3- or 4-fluoro-
phenyl, 2-, 3- or 4-bromophenyl, 2-, 3- or 4-iodophenyl,
5 3,5-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichloro-
phenyl, 2,4-dichlorophenyl, 3,4-difluorophenyl, 3,5-di-
bromophenyl, 2-, 3- or 4-nitrophenyl, 2,4-dinitrophenyl,
2,6-dinitrophenyl, 3,4-dinitrophenyl, 3,5-dinitrophenyl,
2-, 3- or 4-methoxyphenyl, 2-, 3- or 4-ethoxyphenyl, 3-
10 propoxyphenyl, 4-isopropoxyphenyl, 3-butoxyphenyl, 2-
pentyloxyphenyl, 4-tert-butoxyphenyl, 4-hexyloxyphenyl,
3,4-dimethoxyphenyl, 3,4-diethoxyphenyl, 2,5-dimethoxy-
phenyl, 2-, 3- or 4-methylphenyl, 2-, 3- or 4-ethylphenyl,
4-propylphenyl, 3-isopropylphenyl, 2-butylphenyl, 4-
15 hexylphenyl, 3-pentylphenyl, 4-tert-butylphenyl, 3,4-
dimethylphenyl, 2,5-dimethylphenyl, 2-(trifluoromethyl)-
phenyl, 3-(2,3-difluoroethyl)phenyl, 4-(1,1-dichloro-
ethyl)phenyl, 3-(trichloromethyl)phenyl, 2-(dichloromethyl)-
phenyl, 4-(tribromomethyl)phenyl, 3-(2,2,2-trifluoroethyl)-
20 phenyl, 2-(2-chloroethyl)phenyl, 4-(1,2-dichloroethyl)-
phenyl, 3-(3,3,3-trichloropropyl)phenyl, 4-(3-chloro-2-
methylethyl)phenyl, 3-(4-chlorobutyl)phenyl, 2-(3-fluoro-
propyl)phenyl, 3-methyl-4-chlorophenyl, 2-chloro-6-
methylphenyl and 2-methoxy-3-nitrophenyl groups and others
25 can be exemplified.

As to the 1,2,4,6-tetrahydropyridyl-lower alkyl
group which may have, as the substituent, a phenyl group
which may have halogen atoms or lower alkyl groups as

1 the substituents on the phenyl ring, a 1,2,3,6-tetra-
 hydropyridyl group substituted-alkyl group having 1 to 6
 carbon atoms in the alkyl moiety, which may have as the
 substituent, a phenyl group which may have halogen atoms
 5 or alkyl groups having 1 to 6 carbon atoms, as the
 substituents on the phenyl ring, such as 1,2,3,6-tetra-
 hydropyridylmethyl, 2-(1,2,3,6-tetrahydropyridyl)ethyl,
 1-(1,2,3,6-tetrahydropyridyl)ethyl, 3-(1,2,3,6-tetra-
 hydropyridyl)propyl, 4-(1,2,3,6-tetrahydropyridyl)butyl,
 10 1,1-dimethyl-2-(1,2,3,6-tetrahydropyridyl)ethyl, 5-
 (1,2,3,6-tetrahydropyridyl)pentyl, 6-(1,2,3,6-tetra-
 hydropyridyl)hexyl, 2-methyl-3-(1,2,3,6-tetrahydropyridyl)-
 propyl, (4-phenyl-1,2,3,6-tetrahydropyridyl)methyl,
 2-[4-(2-fluorophenyl)-1,2,3,6-tetrahydropyridyl]ethyl,
 15 1-[4-(3-bromophenyl)-1,2,3,6-tetrahydropyridyl]ethyl,
 3-[3-(4-chlorophenyl)-1,2,3,6-tetrahydropyridyl]propyl,
 [4-(4-chlorophenyl)-1,2,3,6-tetrahydropyridyl]methyl,
 [4-(4-methylphenyl)-1,2,3,6-tetrahydropyridyl]methyl,
 [4-(2-(2-ethylphenyl)-1,2,3,6-tetrahydropyridyl]methyl,
 20 1,1-dimethyl-2-[3-(3-propylphenyl)-1,2,3,6-tetrahydro-
 pyridyl]ethyl, 5-[2-(2-tert-butylphenyl)-1,2,3,6-
 tetrahydropyridyl]pentyl, 6-[3-(4-pentylphenyl)-1,2,3,6-
 tetrahydropyridyl]hexyl, and 2-methyl-3-[4-(3-hexylphenyl)-
 1,2,3,6-tetrahydropyridyl]propyl groups and others can
 25 be exemplified.

As to the straight-chain or branched-chain
 unsaturated hydrocarbon residual group which may have or
 may not have an oxygen atom, a sulfur atom, a group of

1 the formula $-N-R^7$ (wherein R^7 is a lower alkyl group) or
a group of the formula $-N \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} N-$ in the unsaturated
hydrocarbon residual group, a straight-chain or branched-
chain unsaturated hydrocarbon residual group having 2 to
5 6 carbon atoms in the unsaturated hydrocarbon residual
moiety, having 1 to 3 double bonds and/or triple bonds
therein, said unsaturated hydrocarbon residual group
may have or may not have an oxygen atom, a sulfur atom,
a group of the formula $-N-R^7$ (wherein R^7 is an alkyl
10 group having 1 to 6 carbon atoms), or a group of the
formula $-N \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} N-$, such as vinylene, 1-propenylene, 1-
methyl-1-propenylene, 2-methyl-1-propenylene, 2-propeny-
lene, 2-butenylene, 1-butenylene, 3-butenylene, 2-
pentenylene, 1-pentenylene, 3-pentenylene, 4-pentenylene,
15 1,3-butadienylene, 1,3-pentadienylene, 2-penten-4-ylene,
2-hexenylene, 1-hexenylene, 5-hexenylene, 3-hexenylene,
4-hexenylene, 3,3-dimethyl-1-propenylene, 2-ethyl-1-
propenylene, ethynylene, 2-propynylene, 1-propynylene,
1,1-dimethyl-2-propynylene, 3,3-dimethyl-1-propynylene,
20 2-butynylene, 3-butynylene, 1-butynylene, 2-pentynylene,
1-pentynylene, 3-pentynylene, 4-pentynylene, 2-hexynylene,
1-hexynylene, 3-hexynylene, 4-hexynylene, 5-hexynylene,
1,3-hexadienylene, 1,4-hexadienylene, 1,3,5-hexatrienylene,
1-propenylenoxy, 1-methyl-1-propenylenoxy, 2-methyl-1-
25 propenylenoxy, 2-propenylenoxy, 2-butenylenoxy, 1-
propenylenoxymethylene, 1-butenylenoxy, 3-butenylenoxy,
2-pentenyleneoxy, 1-pentenyleneoxy, 3-pentenyleneoxy,
4-pentenyleneoxy, 1,3-butadienylenoxy, 1,3-pentadienylenoxy,

- 1 oxy, 2-penten-4-yleneoxy, 2-hexenyleneoxy, 1-hexenyleneoxy,
5-hexenyleneoxy, 3-hexenyleneoxy, 4-hexenyleneoxy, 3,3-
dimethyl-1-propenyleneoxy, 2-ethyl-1-propenyleneoxy,
ethynyleneoxy, 2-propynyleneoxy, 1-propynyleneoxy, 1,1-
5 dimethyl-2-propynyleneoxy, 3,3-dimethyl-1-propynyleneoxy,
3-butyneyleneoxy, 1-butyneyleneoxy, 2-pentyneyleneoxy, 1-
pentyneyleneoxy, 3-pentyneyleneoxy, 4-pentyneyleneoxy, 2-
hexynyleneoxy, 1-hexynyleneoxy, 3-hexynyleneoxy, 4-
hexynyleneoxy, 5-hexynyleneoxy, 1,3-hexadienyleneoxy, 1,4-
10 hexadienyleneoxy, 1,3,5-hexatrienyleneoxy, 1-propenylene-
oxyethylene, 1-propenyleneoxypropylene, 1-methyl-1-
propenyleneoxymethylene, 2-methyl-1-propenyleneoxyethylene,
2-propenyleneoxypropylene, 2-butenyleneoxymethylene,
1-butenyleneoxyethylene, 2-pentenyleneoxymethylene,
15 1,3-butadienyleneoxymethylene, 1,3-pentadienyleneoxy-
methylene, 1-propynyleneoxymethylene, 2-propynylene-
oxyethylene, ethynyleneoxymethylene, 3-butyneyleneoxy-
methylene, 1-butyneyleneoxyethylene, 1-pentyneyleneoxy-
methylene, 3,3-dimethyl-1-propynyleneoxymethylene, 1-
20 propenyleneethio, 1-methyl-1-propenyleneethio, 2-methyl-
1-propenyleneethio, 2-propenyleneethio, 2-butenyleneethio,
1-propenyleneethiomethylene, 1-butenyleneethio, 3-butenylene-
ethio, 2-pentenyleneethio, 1-pentenyleneethio, 3-pentenylene-
ethio, 4-pentenyleneethio, 1,3-butadienyleneethio, 1,3-
25 pentadienyleneethio, 2-penten-4-ylenethio, 2-hexenylene-
ethio, 1-hexenyleneethio, 5-hexenyleneethio, 3-hexenylene-
ethio, 4-hexenyleneethio, 3,3-dimethyl-1-propenyleneethio,
2-ethyl-1-propenyleneethio, ethynyleneethio, 2-propynyleneethio,

- 1 1-propynylene-thio, 1,1-dimethyl-2-propynylene-thio, 3,3-dimethyl-1-propynylene-thio, 3-butyne-thio, 1-butyne-thio, 2-pentyne-thio, 1-pentyne-thio, 3-pentyne-thio, 4-pentyne-thio, 2-hexyne-thio, 1-hexyne-thio, 3-hexyne-thio, 4-hexyne-thio, 5-hexyne-thio, 1,3-hexadiene-thio, 1,4-hexadiene-thio, 1,3,5-hexatriene-thio, 1-propenylene-thio-ethylene, 1-propenylene-thio-propylene, 1-methyl-1-propenylene-thio-methylene, 2-methyl-1-propenylene-thio-ethylene, 2-propenylene-thio-propylene, 2-butenylene-thio-methylene, 1-butenylene-thio-ethylene, 2-pentenylene-thio-methylene, 1,3-butadiene-thio-methylene, 1,3-pentadiene-thio-methylene, 1-propynylene-thio-methylene, 2-propynylene-thio-ethylene, ethynylene-thio-methylene, 3-butyne-thio-methylene, 1-butyne-thio-ethylene, 1-pentyne-thio-methylene, 3,3-dimethyl-1-propynylene-thio-methylene, 3,3-dimethyl-1-propynylene-thio-methylene, N-methyl-N-(1-propenylene)amino, N-ethyl-N-(1-methyl-1-propenylene)amino, N-propyl-N-(2-methyl-1-propenylene)amino, N-n-butyl-N-(2-propenylene)amino, N-pentyl-N-(2-butenylene)amino, N-methyl-N-(1-propenylene)aminomethylene, N-hexyl-N-(1-butenylene)amino, N-methyl-N-(3-butenylene)amino, N-ethyl-N-(2-pentenylene)amino, N-propyl-N-(1-pentenylene)amino, N-tert-butyl-N-(3-pentenylene)amino, N-pentyl-N-(4-pentenylene)amino, N-hexyl-N-(1,3-butadienylene)amino, N-methyl-N-(1,3-pentadienylene)amino, N-ethyl-N-(2-penten-4-ylene)amino, N-propyl-N-(2-hexenylene)amino, N-n-butyl-N-(1-hexenylene)amino, N-pentyl-N-(5-hexenylene)-

- 1 amino, N-hexyl-N-(3-hexenylene)amino, N-methyl-N-(1-propynylene)amino, N-ethyl-N-(1,1-dimethyl-2-propynylene)amino, N-propyl-N-(3,3-dimethyl-1-propynylene)amino, N-tert-butyl-N-(3-butyne)amino, N-pentyl-N-(1-butyne)amino, N-hexyl-N-(2-pentyne)amino, N-methyl-N-(1-pentyne)amino, N-ethyl-N-(3-pentyne)amino, N-propyl-N-(4-pentyne)amino, N-butyl-N-(2-hexynylene)amino, N-pentyl-N-(1-hexynylene)amino, N-hexyl-N-(3-hexynylene)amino, N-methyl-N-(4-hexynylene)amino, N-ethyl-N-(5-hexynylene)amino, N-propyl-N-(1,3-hexadienylene)amino, N-tert-butyl-N-(1,4-hexadienylene)amino, N-pentyl-N-(1,3,5-hexatrienylene)amino, N-hexyl-N-(1-propenylene)aminoethylene, N-methyl-N-(1-propenylene)aminopropylene, N-ethyl-N-(1-methyl-1-propenylene)aminomethylene, N-propyl-N-(2-methyl-1-propenylene)aminoethylene, N-butyl-N-(2-propenylene)aminopropylene, N-pentyl-N-(2-butenylene)aminomethylene, N-hexyl-N-(1-butenylene)aminoethylene, N-methyl-N-(2-pentenylene)aminomethylene, N-ethyl-N-(1,3-butadienylene)aminoethylene, N-propyl-N-(1,3-pentadienylene)aminomethylene, N-methyl-N-(1-propynylene)aminomethylene, N-butyl-N-(2-propynylene)aminoethylene, N-pentyl-N-ethynyleneaminomethylene, N-hexyl-N-(3-butyne)aminomethylene, N-methyl-N-(1-butyne)aminoethylene, N-ethyl-N-(1-pentyne)aminomethylene, N-methyl-N-(3,3-dimethyl-1-propynylene)aminomethylene, 4-(1-propenylene)-1-piperazinyl, 4-(1-methyl-1-propenylene)-1-piperazinyl, 4-(2-methyl-1-propenylene)-1-piperazinyl, 4-(2-propenylene)-1-piperazinyl,

- 1 4-(2-butenylene)-1-piperaziny1, 4-(1-propenylene)-1-piperaziny1methylene, 4-(1-butenylene)-1-piperaziny1, 4-(3-butenylene)-1-piperaziny1, 4-(2-pentenylene)-1-piperaziny1, 4-(1-pentenylene)-1-piperaziny1, 4-(3-pentenylene)-1-piperaziny1, 4-(4-pentenylene)-1-piperaziny1, 4-(1,3-butadienylene)-1-piperaziny1, 4-(1,3-pentadienylene)-1-piperaziny1, 4-(2-penten-4-ynylene)-1-piperaziny1, 4-(2-hexenylene)-1-piperaziny1, 4-(1-hexenylene)-1-piperaziny1, 4-(5-hexenylene)-1-piperaziny1, 4-(1-butynylene)-1-piperaziny1ethylene, 4-(1-pentynylene)-1-piperaziny1methylene, 4-(3,3-dimethyl-1-propynylene)-1-piperaziny1methylene, 4-(3-hexenylene)-1-piperaziny1, 4-(4-hexenylene)-1-piperaziny1, 4-(3,3-dimethyl-1-propenylene)-1-piperaziny1, 4-(2-ethyl-1-propenylene)-1-piperaziny1, 4-(ethynylene)-1-piperaziny1, 4-(2-propynylene)-1-piperaziny1, 4-(1-propynylene)-1-piperaziny1, 4-(1,1-dimethyl-2-propynylene)-1-piperaziny1, 4-(3,3-dimethyl-1-propynylene)-1-piperaziny1, 4-(3-butynylene)-1-piperaziny1, 4-(1-butynylene)-1-piperaziny1, 4-(2-pentynylene)-1-piperaziny1, 4-(1-pentynylene)-1-piperaziny1, 4-(3-pentylene)-1-piperaziny1, 4-(4-pentynylene)-1-piperaziny1, 4-(2-hexynylene)-1-piperaziny1, 4-(1-hexynylene)-1-piperaziny1, 4-(3-hexynylene)-1-piperaziny1, 4-(4-hexynylene)-1-piperaziny1, 4-(5-hexynylene)-1-piperaziny1, 4-(1,3-hexadienylene)-1-piperaziny1, 4-(1,4-hexadienylene)-1-piperaziny1, 4-(1,3,5-hexatrienylene)-1-piperaziny1, 4-(1-propenylene)-1-piperaziny1ethylene, 4-(1-propenylene)-1-piperaziny1propylene, 4-(1-methyl-1-

- 1 propenylene)-1-piperazinylmethylene, 4-(2-methyl-1-propenylene)-1-piperazinylethylene, 4-(2-propenylene)-1-piperazinylpropylene, 4-(2-butenylene)-1-piperazinylmethylene, 4-(1-butenylene)-1-piperazinylethylene, 4-(2-pentenylene)-1-piperazinylmethylene, 4-(1,3-butadienylene)-1-piperazinylmethylene, 4-(1,3-pentadienylene)-1-piperazinylmethylene, 4-(1-propynylene)-1-piperazinylmethylene, 4-(2-propynylene)-1-piperazinylethylene, 4-ethylene-1-piperazinylmethylene and 4-(3-butyne)-1-piperazinylmethylene groups and others can be exemplified.

- As to the tetrazolyl group which may have a lower alkyl group as the substituent, a tetrazolyl group which may have an alkyl group having 1 to 6 carbon atom as the substituent, such as tetrazolyl, 1-methyl-5-tetrazolyl, 1-ethyl-5-tetrazolyl, 1-propyl-5-tetrazolyl, 1-tert-butyl-5-tetrazolyl, 1-pentyl-5-tetrazolyl, 1-hexyl-5-tetrazolyl, 5-methyl-1-tetrazolyl, 5-isopropyl-1-tetrazolyl, 5-n-butyl-1-tetrazolyl, and 5-hexyl-1-tetrazolyl groups and others can be exemplified.

As to the phenyl group which may have a hydroxyl group as the substituent, phenyl, 2-, 3- or 4-hydroxyphenyl group can be exemplified.

- As to the straight-chain or branched-chain unsaturated hydrocarbon group which may have or may not have an oxygen atom or a group of the formula -N-R^7 (wherein R^7 is a lower alkyl group), a straight-chain or branched-chain unsaturated hydrocarbon residual group

- 1 having 2 to 6 carbon atoms in the unsaturated hydro-
carbon residual moiety, having 1 to 3 double bonds and/or
triple bonds therein, said unsaturated hydrocarbon
residual group may have or may not have an oxygen atom or
5 a group of the formula -N-R^7 (wherein R^7 is an alkyl
group having 1 to 6 carbon atoms), such as vinylene,
1-propenylene, 1-methyl-1-propenylene, 2-methyl-1-
propenylene, 2-propenylene, 2-butenylene, 1-butenylene,
3-butenylene, 2-pentenylene, 1-pentenylene, 3-pentenylene,
10 4-pentenylene, 1,3-butadienylene, 1,3-pentadienylene,
2-penten-4-ynylene, 2-hexenylene, 1-hexenylene, 5-
hexenylene, 3-hexenylene, 4-hexenylene, 3,3-dimethyl-1-
propenylene, 2-ethyl-1-propenylene, ethynylene, 2-
propynylene, 1-propynylene, 1,1-dimethyl-2-propynylene,
15 3,3-dimethyl-1-propynylene, 2-butyne, 3-butyne,
1-butyne, 2-pentyne, 1-pentyne, 3-pentyne,
4-pentyne, 2-hexynylene, 1-hexynylene, 3-hexynylene,
4-hexynylene, 5-hexynylene, 1,3-hexadienylene, 1,4-
hexadienylene, 1,3,5-hexatrienylene, 1-propenyleneoxy,
20 1-methyl-1-propenyleneoxy, 2-methyl-1-propenyleneoxy,
2-propenyleneoxy, 2-butenyleneoxy, 1-propenyleneoxy
methylene, 1-butenyleneoxy, 3-butenyleneoxy, 2-penten-
yleneoxy, 1-pentenyleneoxy, 3-pentenyleneoxy, 4-penten-
yleneoxy, 1,3-butadienyleneoxy, 1,3-pentadienyleneoxy,
25 2-penten-4-ynyleneoxy, 2-hexenyleneoxy, 1-hexenyleneoxy,
5-hexenyleneoxy, 3-hexenyleneoxy, 4-hexenyleneoxy, 3,3-
dimethyl-1-propenyleneoxy, 2-ethyl-1-propenyleneoxy,
ethynyleneoxy, 2-propynyleneoxy, 1-propynyleneoxy, 1,1-


- 1 dimethyl-2-propynyleneoxy, 3,3-dimethyl-1-propynyleneoxy,
3-butynyleneoxy, 1-butynyleneoxy, 2-pentynyleneoxy, 1-
pentynyleneoxy, 3-pentynyleneoxy, 4-pentynyleneoxy,
2-hexynyleneoxy, 1-hexynyleneoxy, 3-hexynyleneoxy, 4-
5 hexynyleneoxy, 5-hexynyleneoxy, 1,3-hexadienyleneoxy,
1,4-hexadienyleneoxy, 1,3,5-hexatrienyleneoxy, 1-
propenyleneoxyethylene, 1-propenyleneoxypropylene, 1-
methyl-1-propenyleneoxymethylene, 2-methyl-1-propenylene-
oxyethylene, 2-propenyleneoxypropylene, 2-butenyleneoxy-
10 methylene, 1-butenyleneoxyethylene, 2-pentenyleneoxy-
methylene, 1,3-butadienyleneoxymethylene, 1,3-pentadieny-
leneoxymethylene, 1-propynyleneoxymethylene, 2-propyny-
leneoxyethylene, ethynyleneoxymethylene, 3-butynylene-
oxymethylene, 1-butynyleneoxyethylene, 1-pentynylene-
15 oxymethylene, 3,3-dimethyl-1-propynyleneoxymethylene,
N-methyl-N-(1-propenylene)amino, N-ethyl-N-(1-methyl-1-
propenylene)amino, N-propyl-N-(2-methyl-1-propenylene)-
amino, N-n-butyl-N-(2-propenylene)amino, N-pentyl-N-
(2-butenylene)amino, N-methyl-N-(1-propenylene)amino-
20 methylene, N-hexyl-N-(1-butenylene)amino, N-methyl-N-
(3-butenylene)amino, N-ethyl-N-(2-pentenylene)amino,
N-propyl-N-(1-pentenylene)amino, N-tert-butyl-N-(3-
pentenylene)amino, N-pentyl-N-(4-pentenylene)amino, N-
hexyl-N-(1,3-butadienylene)amino, N-methyl-N-(1,3-
25 pentadienylene)amino, N-ethyl-N-(2-penten-4-ynylene)-
amino, N-propyl-N-(2-hexenylene)amino, N-n-butyl-N-
(1-hexenylene)amino, N-pentyl-N-(5-hexenylene)amino,
N-hexyl-N-(3-hexenylene)amino, N-methyl-N-(1-propynylene)-

- 1 amino, N-ethyl-N-(1,1-dimethyl-2-propynylene)amino,
N-propyl-N-(3,3-dimethyl-1-propynylene)amino, N-tert-
butyl-N-(3-butynylene)amino, N-pentyl-N-(1-butynylene)-
amino, N-hexyl-N-(2-pentynylene)amino, N-ethyl-N-(3-
5 pentynylene)amino, N-methyl-N-(1-pentynylene)amino,
N-propyl-N-(4-pentynylene)amino, N-butyl-N-(2-hexynylene)-
amino, N-pentyl-N-(1-hexynylene)amino, N-hexyl-N-(3-
hexynylene)amino, N-methyl-(4-hexynylene)amino, N-ethyl-
N-(5-hexynylene)amino, N-propyl-N-(1,3-hexadienylene)-
10 amino, N-tert-butyl-N-(1,4-hexadienylene)amino, N-
pentyl-N-(1,3,5-hexatrienylene)amino, N-hexyl-N-(1-
propenylene)aminoethylene, N-methyl-N-(1-propenylene)-
aminopropylene, N-ethyl-N-(1-methyl-1-propenylene)-
aminomethylene, N-propyl-N-(2-methyl-1-propenylene)-
15 aminoethylene, N-butyl-N-(2-propenylene)aminopropylene,
N-pentyl-N-(2-butenylene)aminomethylene, N-hexyl-N-
(1-butenylene)aminoethylene, N-methyl-N-(2-pentenylene)-
aminomethylene, N-ethyl-N-(1,3-butadienylene)aminomethylene,
N-propyl-N-(1,3-pentadienylene)aminomethylene, N-methyl-
20 N-(1-propynylene)aminomethylene, N-butyl-N-(2-propynylene)-
aminoethylene, N-pentyl-N-ethynyleneaminomethylene,
N-hexyl-N-(3-butynylene)aminomethylene, N-methyl-N-(1-
butynylene)aminoethylene, N-ethyl-N-(1-pentynylene)-
aminomethylene and N-methyl-N-(3,3-dimethyl-1-propyny-
25 lene)aminomethylene groups and others can be exemplified.

Dihydropyridine derivatives and salts thereof represented by the general formula (1) according to the present invention can be prepared by various methods,

1 for examples the derivatives can be prepared by the following Reaction process formula-1.

Reaction process formula-1



(wherein X is a hydroxyl group or a halogen atom; and
 5 R^1 , R^2 , R^3 , R^4 and R^5 are the same as defined above).

In the reaction of compound (2) with compound
 (3), when X is a hydroxyl group, reaction conditions
 usually employed in esterification reaction can be used.
 The reaction may be conducted generally in the presence
 10 of a catalyst which is usually used in esterification
 reactions. As to the typical catalysts, there can be
 exemplified inorganic acids, such as hydrogen chloride,
 concentrated sulfuric acid, phosphoric acid, poly-
 phosphoric acids, boron trifluoride and perchloric acid
 15 and others; organic acids, such as trifluoroacetic acid,
 trifluoromethanesulfonic acid, naphthalenesulfonic
 acids, p-toluenesulfonic acid, benzenesulfonic acid and
 ethanesulfonic acid and others; and dehydrating agents,
 such as trifluoromethanesulfonic acid anhydride, thionyl
 20 chloride, tetramethylurea oxalyl chloride, acetone

1 dimethyl acetal, dicyclohexylcarbodiimide (DCC), 1-
alkyl-2-halogenopyridinium halide or tosylate, N,N-
carbonyldiimidazol and others can be exemplified.
Additionally, acidic ion-exchange resins can also be
5 used as the catalysts. The amount of these catalysts is
not restricted in the specific range, and they can be
used in any amount usually used in common esterification
reaction.

The reaction may be carried out in the absence
10 or presence of a solvent. As to the solvent used in
the reaction, any solvent usually used in common
esterification reaction may effectively be used.
Specifically, as to the solvents, aromatic hydrocarbons,
for example benzene, toluene and xylene; halogenated
15 hydrocarbons, for example dichloromethane, dichloro-
ethane, chloroform, carbon tetrachloride and others;
ethers, for example diethyl ether, tetrahydrofuran,
dioxane, ethylene glycol monomethyl ether, pyridine and
others, and mixed solvents thereof can be exemplified.

20 In the above-mentioned reaction, the ratio of
the amount of compound (3) to the amount of compound (2)
can be selected from a wide range, and the former is
used in an equimolar quantity to 5 times the molar quantity,
preferably, the equimolar quantity to 2 times the molar
25 quantity to the latter.

In carrying out of the above-mentioned reaction,
the yield of the objective product can be increased by
removing the water formed in the reaction from the


1 reaction system by using a dehydrating agent, such as
anhydrous calcium chloride, anhydrous copper sulfate,
anhydrous calcium sulfate, phosphorus pentoxide or the
like. The reaction temperature of the reaction may be
5 selected optionally, and there is not any restriction to
the temperature, generally, the reaction may be carried
out in the range from about -20° to 200°C , preferably,
from about 0°C to 150°C . The reaction is completed,
generally in about 10 minutes to 20 hours depend on
10 the kind of the starting materials and the reaction
condition.

In the above-mentioned reaction, when X is a
halogen atom, the objective product can be obtained by
carrying out the reaction under conditions of dehydro-
15 halogenating reaction. The dehydrohalogenating reaction
is carried out by using a basic compound as the dehydro-
halogenating agent. As to the basic compound, any known
basic compound can be used, for example inorganic basic
compounds such as sodium hydroxide, potassium hydroxide,
20 sodium carbonate, potassium carbonate, sodium hydrogen
carbonate, potassium hydrogen carbonate, silver carbo-
nate and others; alcoholates such as sodium methylete,
sodium ethylete and others; organic basic compounds
such as triethylamine, pyridine, N,N-dimethylaniline
25 and others can be exemplified.

The dehydrohalogenating reaction can advan-
tageously be carried out in the presence of a solvent,
and any inert solvent which does not give any adverse

1 effect to the reaction can be used. As to the solvents,
alcohols, such as methanol, ethanol, propanol, butanol,
ethylene glycol and others; ethers, such as dimethyl
ether, tetrahydrofuran, dioxane, ethylene glycol dimethyl
5 ether, diethylene glycol dimethyl ether, and others;
ketones such as acetone, methyl ether ketone and others;
aromatic hydrocarbons, such as benzene, toluene, xylene
and others; esters, such as methyl acetate, ethyl
acetate and others; aprotic polar solvents, such as
10 N,N-dimethylformamide, dimethyl sulfoxide, hexamethyl-
phosphoryl triamide and others can be exemplified. Said
reaction can also be carried out in the presence of a
metal iodide, such as sodium iodide, potassium iodide
and others. The ratio of the amount of a compound (3)
15 to the amount of a compound (2) is not specifically
restricted and can be selected from a wide range,
generally an equimolar quantity to 5 times the molar
quantity, preferably an equimolar quantity to 2 times
the molar quantity of the former is used to the latter.
20 The reaction temperature is also not specifically
restricted, and generally the reaction is carried out at
a room temperature to 200°C, preferably at from a room
temperature to 160°C. The reaction is preferably carried
out in 1 to 30 hours. Thus, dihydropyridine derivatives
25 represented by the general formula (1) can be prepared.

1 Reaction process formula-2



(wherein R^1 , R^2 , R^3 , R^4 and R^5 are the same as defined above).

The reaction of compound (4) with compound (5) in the above-mentioned reaction process formula-2 can also be carried out in a suitable solvent in the presence or absence of a catalyst.

As to the solvent, alcohols, such as methanol, ethanol, propanol, isopropanol, butanol, ethylene glycol, and others; ethers, such as tetrahydrofuran, dioxane, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether and others; aromatic hydrocarbons, such as benzene, toluene, xylene and others; halogenated hydrocarbons, such as methylene chloride, chloroform, 1,2-dichloromethane, and others; aprotic polar solvents, such as dimethyl sulfoxide, N,N-dimethylformamide, hexamethylphosphoryl triamide and others; carboxylic acids,

1 such as acetic acid, propionic acid and others; and
pyridine can be exemplified.

As to the catalyst used in the reaction, organic
basic compounds, such as pyridine, piperidine, triethylamine,
5 diethylamine, 1,8-diazabicyclo[5,4,0]undecene-5 (DBU)
and others; metal alcoholates, such as sodium ethylate,
sodium methylate and others, inorganic basic compounds,
such as sodium hydroxide, potassium hydroxide, potassium
carbonate, potassium acetate and others; mineral acids,
10 such as hydrochloric acid, sulfuric acid and others;
carboxylic acid, such as acetic acid, propionic acid
and others, Lewis acids, such as boron trifluoride can
be exemplified.

As to the ratio of the amount of compound (4)
15 to the amount of compound (5), the latter may be used in
an equimolar quantity, preferably an equimolar quantity
to 2 times the molar quantity to the former. As to the
amount of the catalyst, 0.01 to 10 times the molar
quantity, preferably, 0.1 to 5 times the molar quantity
20 of the catalyst may be used to compound (4). The reaction
may be carried out, generally at -20 to 200°C, preferably,
-20 to 150°C, and the reaction is completed generally,
in 10 minutes to 50 hours.

The reaction of compound (6) with compound (7)
25 can advantageously be carried out in the presence of a
solvent. As to the solvent, any inert solvent which does
not give any adverse effect to the reaction can be used,
for example, ketones, such as acetone; halogenated

1 hydrocarbons such as chloroform and others; alcohols,
such as methanol, ethanol, propanol, isopropanol, ethylene
glycol and others; ethers, such as diethyl ether,
tetrahydrofuran, dioxane, ethylene glycol dimethyl ether,
5 diethylene glycol dimethyl ether and others; aromatic
hydrocarbons, such as benzene, toluene, xylene and others;
esters, such as methyl acetate, ethyl acetate and others;
carboxylic acid, such as acetic acid, propionic acid
and others; organic basic compounds such as pyridine and
10 others; and aprotic polar solvent such as N,N-dimethyl-
formamide, dimethyl sulfoxide, hexamethylphosphoryl
triamide and others can be exemplified.

The ratio of the amount of compound (6) to the
amount of compound (7) may be an equimolar quantity to
15 10 times the molar quantity, preferably an equimolar
quantity to 2 times the molar quantity ^{of the latter may be used} to the former.
The reaction may generally be carried out at -20° to 200°C,
preferably at 50° to 150°C. The reaction is generally
completed in 10 minutes to 20 hours, then the desired
20 compound represented by formula (1) can be obtained

In conducting the reaction of compound (4)
with compound (5) to form compound (6), then reacting
compound (6) with compound (7) to prepare the desired
compound (1), the intermediate compound (6) may not be
25 separated from the reaction system, thus compound (5)
and compound (7) may be existed within the same reaction
system and make them reacted in a simultaneous (in one
step) reaction.

1 Among the dihydropyridine derivatives represented
by the general formula (1), those having, as for the
symbol R^8 , a phenyl group which contains at least one
hydroxyl group as the substituent may be prepared by
5 hydrolyzing a compound among those represented by the
general formula (1) having, as for the symbol R^8 , a
phenyl group which contains at least one substituent
selected from the group consisting of a lower alkoxy
group, a tetrahydropyranyloxy group, a lower alkanoyloxy
10 group and a lower alkoxy-lower alkoxy group.

The hydrolyzing reaction of the compound (1)
[hereinafter referred to as "compound (1a)"] having, as
for the symbol R^8 , a phenyl group which contains at
least one substituent selected from the group consisting
15 of a lower alkoxy group, a tetrahydropyranyloxy group and
a lower alkoxy-lower alkoxy group is carried out without
in a solvent or with in a suitable solvent, by reacting
an acid. As to the solvent used in this hydrolyzing
reaction, water; nitrobenzene; aromatic hydrocarbons such
20 as benzene, toluene, xylene and others; saturated,
hydrocarbons such as hexane, octane and others; lower
alcohols such as methanol, ethanol, isopropanol and
others; ethers such as dioxane, tetrahydrofuran and
others; ketones such as acetone and others; acetic acid;
25 acetonitrile and mixed solvents thereof can be exemplified.
As to the acid used in this hydrolyzing reaction, mineral
acids such as hydrochloric acid, hydrobromic acid, sulfuric
acid and others; p-toluenesulfonic acid; pyridine

1 p-toluenesulfonate; carboxylic acids such as acetic
acid, propionic acid and others; aluminium chloride; tin
chloride; boron trifluoride; zinc chloride and others
can be exemplified. The amount of the acid to be
5 used to the amount of compound (1a) may be at least an
equimolar quantity, generally a large excess quantity
may be used. The reaction may be carried out generally
from -30° to 200°C, preferably from -30° to 100°C, and
the reaction is generally completed in about 0.5 to 8
10 hours.

The hydrolyzing reaction of the compound (1)
having, as for the symbol R^8 , a phenyl group which
contains at least one lower alkanoyloxy group, is carried
out under reaction conditions widely employed in hydro-
15 lyzing reaction of esters. For example, the hydrolyzing
reaction is carried out under conditions of in the
presence of an acid or alkali catalyst, in an inert
solvent at 0° to 100°C, for 1 to 5 hours. As to the
catalyst, inorganic acids such as hydrochloric acid,
20 sulfuric acid, aluminium chloride and others; organic
acids such as acetic acid, formic acid and others;
inorganic basic compounds such as sodium hydroxide
sodium carbonate, potassium hydroxide and others;
ammonia; organic basic compounds such as triethylamine
25 and others can be exemplified. As to the inert solvent,
water; alcohols such as methyl alcohol, ethyl alcohol
and others; carboxylic acids such as acetic acid, propionic
acid and others; ethers such as diethyl ether and others;

1 amides such as dimethylformamide, acetamide and others
can be exemplified.

Among the dihydropyridine derivatives represented by the general formula (1), those having, as for
5 the symbol R^8 , a phenyl group which contains at least one substituent selected from the group consisting of a lower alkoxy group, a tetrahydropyranyloxy group and a lower alkoxy-lower alkoxy group, can also be prepared by alkylating a compound hereinafter referred to as
10 "compound (1b)" having, as for the symbol R^8 , a phenyl group which contains at least one hydroxyl group as the substituent. Said alkylating reaction is carried out under conditions usually employed in common alkylating reaction. For example, alkylation is carried out by
15 using an alkylating agent in the presence of a basic compound. As to the basic compound used in this reaction, alkali metals such as metallic sodium, metallic potassium and others; and the hydrides, hydroxides, carbonates, bicarbonates or alcoholates of these alkali metals;
20 aromatic amines compounds such as pyridine, piperidine, and others; organic basic compounds such as triethylamine, N,N-diethylaniline, 1,8-diazabicyclo[5,4,0]undecene-7 (DBU) and others can be exemplified. As to the alkylating agent, a lower alkyl halide, a tetrahydropyranyl halide,
25 dihydropyran, a lower alkoxy-lower alkyl halide, a dialkyl sulfate, a diazoalkane and others can be exemplified.

In using a lower alkyl halide, a tetrahydro-

1 pyranil halide or a lower alkoxy-lower alkoxy halide as
the for alkylating agent, the alkylating reaction is
carried out effectively in a suitable solvent. As to
the solvent to be used, water; a lower alcohols such as
5 methanol, ethanol, isopropanol, n-butanol and others;
ethers such as diethyl ether, dioxane, tetrahydrofuran
and others; ketones such as acetone, methyl ethyl ketone
and others; halogenated hydrocarbons such as chloroform,
dichloroethane and others; aromatic hydrocarbons such as
10 nitrobenzene, chlorobenzene, benzene, toluene, xylene
and others; aprotic polar solvents such as N,N-dimethyl-
formamide, dimethyl sulfoxide and others can be exempli-
fied. The amount of the alkylating agent to the amount
of compound (1b), at least an equimolar quantity,
15 preferably an equimolar quantity to 5 times the molar
quantity may be used to the latter. The reaction is
generally carried out at from -20° to 200°C, preferably
at 0° to 100°C, and is completed in about 10 minutes
to 24 hours.

20 In using a dialkyl sulfate as for the alkylating
agent, the alkylation reaction is carried out in an
inert solvent at a room temperature to 150°C. As to
the dialkyl sulfate, dimethyl sulfate, diethyl sulfate and
others can be exemplified. As to the inert solvent,
25 aromatic hydrocarbons such as benzene, toluene and
others; ethers such as dioxane, tetrahydrofuran, diethyl
ether and others can be exemplified.

In using dihydropyran as for the alkylating

1 agent, the alkylating reaction is carried out in the
presence of an acid, in a solvent, and generally at 0°
to 150°C, preferably at 0° to about 100°C, and the
reaction is completed in 0.5 to 10 hours. As to the acid
5 to be used in this case, mineral acids such as hydro-
chloric acid, hydrobromic acid, sulfuric acid and others;
p-toluenesulfonic acid; pyridine p-toluenesulfonate and
others can be exemplified. As to the solvent, lower
alcohols such as methanol, ethanol, isopropanol and
10 others; ethers such as diethyl ether, dioxane, tetra-
hydrofuran and others; aromatic hydrocarbons such
as benzene, toluene and others; saturated hydrocarbons
such as hexane, octane and others; ketones such as
acetone and; acetic acid; acetonitrile and mixtures of
15 these solvents can be exemplified.





The amount of dihydropyran to the amount of
compound (1b), generally an equimolar quantity, preferably
an equimolar to 5 times the molar quantity of the former
may be used to the latter.

20 Among the dihydropyridine derivatives repre-
sented by the general formula (1), those having, as
for the symbol R⁸, a phenyl group which contains at
least one lower alkanoyloxy group as the substituent,
can also be prepared by acylating a compound (1b).
25 Said acylating reaction is carried out by using an acid
halide such as a lower alkanoic acid halide, or an
alkanoic acid anhydride under conventional method. The
reaction by using acid halide is carried out in an inert

1 solvent, and if necessary in the presence of a dehydro-
halogenating agent such as an amine for example tri-
ethylamine, diisopropylethylamine, pyridine, N,N-
diethylaniline and others, and at -50° to 150°C, in 1
5 to 24 hours. In carrying out the acylating reaction
by using acid anhydride, the reaction is conducted in
an inert solvent at a room temperature to 200°C, in 1
to 10 hours. As to the inert solvent used in the above-
mentioned reaction, aromatic hydrocarbons such as
10 nitrobenzene, chlorobenzene and others; amines such as
pyridine, N,N-dimethylaniline and others; ethers such
as diethyl ether, tetrahydrofuran and others; halogenated
hydrocarbons such as dichloromethane, dichloroethane,
chloroform and others can be exemplified. The amount of
15 the acylating agent used to the amount of compound (1b),
generally at least an equimolar quantity, preferably an
equimolar to 5 times the molar quantity of the former may
be used to the latter.

Compounds represented by the general formula
20 (3) as used for the starting material in the above-
mentioned reaction process formula-1, and compounds
represented by the general formula (4) as used for the
starting material in the above-mentioned reaction process
formula-2 contain novel compounds, said compounds
25 represented by the general formulas (3a), (3b), (3c) and
(3d) are prepared by the following reaction process
formulas-3 to -5.

1 Reaction process formula-3



[wherein R^8 is the same as defined above; R^9 is a hydrogen atom or a lower alkyl group; R^{10} is a lower alkyl group; R^{11} is a carboxyl group or a group of the formula $\text{---} \text{P}(\text{OR}^{10'})_2$ O]

5 (wherein $\text{R}^{10'}$ is a lower alkyl group); $\underline{\text{B}}$ and $\underline{\text{D}}$ are each an unsaturated alkylene group; $\underline{\text{m}}$ and $\underline{\text{o}}$ are 0 or 1 respectively; X^1 is a halogen atom; provided that the carbon

1 atom number in the group of the formula $\text{CH}_2 \begin{array}{l} \nearrow (\text{D})\text{O}^- \\ \searrow \text{R}^{11} \end{array}$ should

not exceed 6].

The reaction of compound (8) with compound (9) can be carried out in the presence of a basic compound, in a solvent. As to the basic compound, inorganic basis such as metallic sodium, metallic potassium, sodium hydride, sodium amide, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydrogen carbonate and others; metal alcoholates such as sodium methylate, sodium ethylate and others; organic basis compounds such as pyridine, piperidine, quinoline, triethylamine, N,N-dimethylaniline and others can be exemplified. As to the solvent, any inert solvent which does not give any adverse effect to the reaction can be used, for example, ethers such as diethyl ether, dioxane, tetrahydrofuran, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether and others; aromatic hydrocarbons such as benzene, toluene, xylene and others; aliphatic hydrocarbons n-hexane, heptane, cyclohexane and others, amines such as pyridine, N,N-dimethylaniline and others; aprotic polar solvents such as N,N-dimethylformamide (DMF), dimethyl sulfoxide (DMSO), hexamethylphosphoryl triamide (HMPA) and others can be exemplified. The reaction is generally carried out at 0° to 150°C, preferably from a room temperature to about 120°C, and is generally completed in 0.5 to 15 hours. The amount of compound (8) to the amount of compound (9) is generally




1 an equimolar quantity, preferably an equimolar quantity
to 2 times the molar quantity of the latter is used to
the former.

The reduction of compound (10) is generally
5 carried out by using a hydrogenation reducing agent. As
to the hydrogenation reducing agent, sodium borohydride,
lithium aluminum hydride, aluminum dialkyl hydride such
as aluminum diisobutyl hydride (DIBAL), diborane and
others. The amount of the hydrogenation reducing agent
10 to the amount of compound (10) is that generally 0.1 to
3 times the molar quantity, preferably 0.5 to 2 times the
molar quantity of the former is used to the latter. The
reaction is carried out generally in a suitable solvent,
for example water; a lower alcohols such as methanol,
15 ethanol, isopropanol and others; ethers such as tetra-
hydrofuran, diethyl ether, diethylene glycol dimethyl
ether and others; aromatic hydrocarbons such as benzene,
toluene, xylene and others can be used, at -60° to 50°C,
preferably at -40°C to a room temperature, for about 10
20 minutes to 5 hours. In carrying out of the reaction
by using lithium aluminum hydride, aluminum dialkyl
hydride or diborane as for the reducing agent, an
anhydrous solvent such as diethyl ether, tetrahydrofuran,
diethylene glycol dimethyl ether, benzene, toluene or
25 xylene may preferably be used.

The halogenation reaction of compound (3a)
thus prepared is carried out in a solvent for example,
an ether such as dioxane, tetrahydrofuran or the like; a

1 chlorinated hydrocarbon such as chloroform, methylene
chloride, carbon tetrachloride or the like, or without
a solvent, by reacting a compound (3a) with a halogenat-
ing agent for example a hydrohalic acid such as hydro-
5 chloric acid or hydrobromic acid; N,N-diethyl-1,2,2-
trichlorovinylamide, phosphorus pentachloride, phosphorus
pentabromide, phosphorus oxychloride, thionyl chloride
or the like, at a room temperature to 150°C, preferably
at a room temperature to 80°C, for 1 to 6 hours. The
10 amount of the halogenating agent to the amount of a
compound (3a) is that at least an equimolar quantity,
generally a large excess quantity of the halogenating
agent may be used to the latter.

Reaction process formula-4



15 [wherein R^8 , B , m and X^1 are the same as defined above;




1 l is an integer of 3 to 6; provided that the number of
carbon atoms in a group of the formula $-(B)_m-CH=CH-(CH_2)_{l-1}$
should not be over 6].

The reaction for preparing compound (12) from
5 compound (11) is carried out in the presence of a basic
compound, in an inert solvent for example, an ether such
as dioxane, tetrahydrofuran, ethylene glycol dimethyl
ether or the like; an aromatic hydrocarbon such as
benzene, toluene, xylene or the like; a lower alcohol such
10 as methanol, ethanol, isopropanol or the like; a polar
solvent such as acetonitrile, N,N-dimethylformamide,
dimethyl sulfoxide or the like. As to the basic compound,
inorganic basic compounds such as calcium carbonate,
sodium carbonate, potassium carbonate, sodium hydrogen
15 carbonate, sodium hydroxide, potassium hydroxide, sodium
amide, sodium hydride, potassium hydride, sodium
methylete, sodium ethylete and others; and organic basic
compounds such as triethylamine, pyridine, quinoline,
1,5-dibiazabicyclo[4,3,0]nonene-5 (DBN), 1,8-diazabicyclo-
20 [5,4,0]undecene-7 (DBU), 1,4-diazabicyclo[2,2,2]octane
(DABCO) and others can be exemplified. The reaction
is generally carried out at a room temperature to 200°C,
preferably at 60° to 120°C, and the reaction is generally
completed in 1 to 24 hours.

25 Reduction reaction of a compound (12) is
carried out under the same conditions similar to those
employed in the reduction of a compound (10) in the above-
mentioned reaction process formula-3.

- 1 The reaction of a compound (13) with a compound (14) is carried out in a suitable inert solvent. As to the solvent to be used in the reaction, ethers such as dioxane, tetrahydrofuran, diethyl ether and others;
- 5 halogenated hydrocarbons such as chloroform, methylene chloride, carbon tetrachloride and others can be exemplified. The reaction temperature is generally at 0 to 150°C, preferably at 0° to about 100°C, and generally, the reaction is completed in 10 minutes to 6 hours.
- 10 The amount of the compound (14) to the amount of the compound (13) is that generally at least an equimolar quantity, preferably in a large excess amount of the former is used to the latter.

Reaction process formula-5



- 15 [wherein R^8 , R^{10} , B , X^1 , m and o are the same as defined


1 above; M is a metal such as copper, sodium, lithium,
 potassium; D' is a saturated- or unsaturated-alkylene
 group; provided that the number of the carbon atoms in
 a group of the formula $-(B)_m-C\equiv C-(D')_o$ should not be
 5 exceed over 6].


The reaction of a compound (3d) with a compound
 (15) is carried out in a suitable solvent. As to the
 solvent used in the reaction, ethers such as diethyl
 ether, dioxane, tetrahydrofuran, ethylene glycol dimethyl
 10 ether, diethylene glycol dimethyl ether and others;
 aromatic hydrocarbon such as benzene, toluene, xylene;
 aliphatic hydrocarbons such as n-hexane, heptane, cyclo-
 hexane and others; amines such as triethylamine, pyridine,
 N,N-dimethylaniline and others; aprotic polar solvents
 15 such as N,N-dimethylformamide (DMF), dimethyl sulfoxide
 (DMSO), hexamethylphosphoryl triamide (HMPA) and others
 can be exemplified. The reaction temperature is generally
 0° to 200°C, preferably from a room temperature to at
 about 150°C, and the reaction is generally completed in
 20 0.5 to 10 hours. The amount of the compound (15) to
 the amount of the compound (3d) is that at least an
 equimolar quantity, preferably an equimolar to 1.5 times
 the molar quantity of the former is used to the latter.


The reduction reaction of a compound (16) can
 25 be carried out under the same conditions those employed
 in the reduction of a compound (10) in reaction process
 formula-3.

Compounds represented by the general formulas

- 1 (3a), (3b), (3c) and (3d) can be converted into a compound represented by the general formula (17),



[wherein Z is a group of the formula ,



- 5 l are the same as defined above],
by oxidizing in the presence of a suitable oxidizing agent.

A part of compounds represented by the general formula (17) correspond to compound (8) which is the starting material used in the above-mentioned reaction process formula-3, thus various desired compounds represented by the general formula (3) can be obtained by carrying out the reaction in sequently in the above-mentioned reaction process formula-3 to -5 and -9 and the above-mentioned oxidation reaction. As to the oxidizing agent used in the above-mentioned oxidation reaction,


15 chromium compounds such as potassium chromate, sodium bichromate, chromium trioxide, pyridinium chlorochromate, anhydrous chromium trioxide-dipyridine complex and others; manganese compounds such as manganese dioxide, potassium permanganate and others; lead tetraacetate; periodic acid;

20 dimethyl sulfoxide; amine oxides such as dimethylamine oxide; pyridine-nitroso compounds such as pyridine-p-

- 1 nitro-N,N-dimethylaniline and others can be exemplified.
 As to the solvent used to the reaction, aromatic hydro-
 carbons such as benzene, toluene, xylene and others;
 halogenated hydrocarbons such as methylene chloride,
 5 chloroform, carbon tetrachloride and others; ethers
 such as diethyl ether, dioxane, tetrahydrofuran and
 others, aliphatic hydrocarbons such as hexane, pentane,
 cyclohexane and others, ketones such as acetone, methyl
 ethyl keton and others; lower alcohols such as methanol,
 10 ethanol, isopropanol; water, acetic acid, dimethyl
 sulfoxide and others can be exemplified. The reaction
 can be carried out by using an acid such as sulfuric
 acid or perchloric acid as the catalyst. The reaction is
 carried out generally at 0° to 200°C, preferably at 0°
 15 to about 150°C, is completed generally in 0.5 to 15 hours.

A compound represented by the general formula
 (11) as used for the starting material in the above-
 mentioned reaction is prepared by for example the following
 reaction process formula-7.

20 Reaction process formula-7




[wherein R^8 , B , m , l and X^1 are the same as defined above].

1 The reaction of a compound (18) with a compound
 (19) is carried out in a suitable solvent. As to the
 solvent used in the reaction, ethers such as diethyl
 ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane
 5 and others; aromatic hydrocarbons such as benzene,
 toluene, xylene and others; aliphatic hydrocarbons such
 as hexane, heptane, pentane and others; and mixtures of
 these solvents and others can be exemplified.

 The reaction is carried out generally at -70° to
 10 50°C, preferably -65°C to about a room temperature,
 and is generally completed in 3 to 30 hours. The amount
 of the compound (19) to the amount of the compound (18)
 is that at least 2 times the molar quantity, preferably
 2 to 3 times the molar quantity of the former is used to
 15 the latter.

 Among the compounds represented by the general
 formula (11), those in which the symbol \underline{m} is 0 thus
 compound (11a) may be prepared by the following reaction
 process formula-8.

20 Reaction process formula-8





- 1 (wherein R^8 , X^1 and \underline{L} are the same as defined above;
and X^2 is a halogen atom).

The reaction of a compound (20) with a compound (21) or compound (22) is generally called as Friedel-Crafts reaction, and can be carried out in a suitable solvent in the presence of a Lewis acid. As to the solvent used in the reaction, any solvent generally used in this type of reaction can advantageously be employed, for example carbon disulfide, nitrobenzene, chlorobenzene, dichloroethane, dichloromethane, trichloroethane, tetrachloroethane and others can be exemplified. As to the Lewis acid used in this reaction, those used conventionally in this type of reaction can preferably be used, for example aluminum chloride, zinc chloride, iron chloride, tin chloride, boron trifluoride, boron tribromide, concentrated sulfuric acid and others can be used. The amount of Lewis acid may be determined suitably, and generally 2 to 6 times the molar quantity, preferably 3 to 4 times the molar quantity of a Lewis acid may be used to a compound (20). The amount of the compound (21) or compound (22) to the amount of the compound (20) is that generally at least an equimolar quantity, preferably an equimolar quantity to 3 times the molar quantity of the former is used to the latter.

The reaction temperature can be selected from a wide range, and generally the reaction is carried out at 0° to 120°C, preferably from 0° to 70°C, and the reaction is completed in 0.5 to about 6 hours.

1 Reaction process formula-9



[wherein R^8 , B , D' , m and o are the same as defined above; R^{12} is a hydroxyl group or a lower alkanoyl group; provided that the number of the carbon atoms in a group of the formula $-(B)_m-C\equiv C-(D')_o-CH_2-$ and a group of the formula $-(B)_m-CH=CH-(D')_o-CH_2-$ should not exceed over 6.]

The reduction of a compound (3f) can be carried out by methods according to various reducing reactions known in the art. For example, catalytic reducing method by using palladium black, palladium carbon, platinum oxide, platinum black, Raney nickel, Lindlar catalyst and others as for the reducing catalysts; reducing methods by using sodium borohydride, lithium aluminum hydride and others as for the reducing agents can be employed.

In carrying out the catalytic reduction, the reaction can be conducted by using a conventional solvent for example water, methanol, ethanol, isopropanol, acetic acid, dioxane, tetrahydrofuran and others, and in the presence of the above-mentioned catalyst, under a normal atmospheric pressure to 20 atmospheric pressure, preferably, from a normal atmospheric pressure to 10


1 atmospheric pressure of hydrogen, and generally at -30°C
to 100°C , preferably 0°C to 50°C . The amount of the
catalyst is generally 0.1 to 40% by weight, preferably
from 1 to 20% by weight of the catalyst is used to a
5 compound (3f), and the reaction is generally completed
in 1 to 12 hours.

In carrying out the reaction by using a reducing
agent such as lithium aluminum hydride, an equimolar
quantity to 20 times the molar quantity, preferably 1.5
10 to 3.5 times the molar quantity of the reducing agent is
used to the compound (2f). The reduction reaction is
carried out in a conventional solvent such as diethyl
ether, tetrahydrofuran, dioxane or the like, and generally
at -30° to 100°C , preferably at 0°C to 70°C , and is
15 completed in 30 minutes to about 12 hours. According to
these procedures, a compound represented by the general
formula (3g) can easily be obtained.


Among the compounds represented by the general
formula (3g), those having a lower alkanoyloxy group as
20 for the symbol R^{12} , can be converted into a compound (3g)
having a hydroxy group as for the symbol R^{12} under the
conditions similar to those employed in the hydrolysis
of a compound (1) wherein R^8 is a phenyl group which
contains at least one lower alkanoyloxy group as the
25 substituent.

A part of compounds represented by the general
formula (3f) can be prepared by a method according to the
following reaction process formula-10.

1 Reaction process formula-10



(3d)




(3h)

[wherein R^8 , B , m , X^1 , M , D' and o are the same as defined above; R^{13} is a hydroxyl group, tetrahydropyranyloxy group, a lower alkoxy-lower alkoxy group or a lower alkanoyl group; provided that the number of the carbon atoms in a group of the formula $-(B)_m-C\equiv C-(D')_o-CH_2-$ should not exceed over 6].

The reaction of a compound (3d) with a compound (23) can be carried out under the same reaction conditions employed in the reaction of a compound (3d) with a compound (15) in the above-mentioned reaction process formula-5.

Reaction process formula-11



(8a)


(3h)

[wherein R^8 , B , and m are the same as defined above].

The reduction reaction of a compound (8a) can

- 1 be carried out under the same reaction conditions employed in the reduction of a compound (10) in the above-mentioned reaction process formula-3.

Reaction process formula-12



- 5 [wherein R^8 , D, D' and O are the same as defined above; Y is an oxygen atom, a sulfur atom, a group of the formula $\text{-N} \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} \text{N-}$ or a group of the formula -N-R^7 (wherein R^7 is the same as defined above); provided that the number of the carbon atoms in a group of the formula $\text{-(D)}_{\text{O}} - \text{CH}_2 - \text{Y} - (\text{D}')_{\text{O}} - \text{CH}_2 -$ should not be exceeded over 6].
- 10

The reaction of a compound (24) with a compound (25) can be carried out in a suitable solvent or without solvent, in the absence or presence of a basic compound. As to the solvent used in the reaction, water; lower


15 alcohols such as methanol, ethanol, isopropanol and others; halogenated hydrocarbons such as dichloromethane, dichloroethane, chloroform, carbon tetrachloride and others; ethers such as diethyl ether, tetrahydrofuran, dioxane and others; aliphatic hydrocarbons such as n-

1 hexane, octane, cyclohexane and others; aromatic hydro-
carbons such as benzene, toluene, xylene and others;
aprotic polar solvents such as acetone, acetonitrile,
N,N-dimethylformamide, dimethyl sulfoxide, hexamethyl-
5 phosphoryl triamide and others; and mixtures of these
solvents can be exemplified. As to the solvent used in
the reaction, inorganic basic compounds such as sodium
hydroxide, potassium hydroxide, sodium carbonate,
potassium carbonate, sodium hydrogen carbonate, potassium
10 hydrogen carbonate, sodium amide, sodium hydride and
others; alcoholates such as sodium methylate, sodium
ethylate and others; organic basic compounds such as
triethylamine, pyridine, N,N-dimethylaniline and others
can be exemplified.

15 The said reaction can be advantageously proceed
by using a charge-transfer catalyst such as tetrabutyl-
ammonium bromide and others. The reaction is carried
out generally, at 0° to 150°C, preferably at 0° to 120°C,
and is completed in about 1 to 10 hours. The ratio of
20 the amount of a compound (24) to the amount of a compound
(25) is that generally an equimolar quantity, preferably
an equimolar quantity to 10 times the molar quantity of
the former is used to the latter.

A compound (4) used as the starting material
25 in the reaction process formula-2 can be prepared by
methods according to reaction process formula-13 and -14
as follows.

1 Reaction process formula-13




[wherein R⁵ is the same as defined above].

The reaction of a compound (26) with a compound (27) in the above-mentioned reaction process formula-13 is carried out in a suitable solvent in the presence of a catalyst. As to the catalyst, basic compounds for example, organic basic compounds such as triethylamine, pyridine, N,N-dimethylaniline and others; inorganic basic compounds such as sodium acetate, potassium carbonate; and acidic compounds for example, sulfonic acids such as p-toluenesulfonic acid and others; Lewis acids such as boron trifluoride and others can be exemplified. As to the solvents, aromatic hydrocarbons such as benzene, toluene, xylene and others; esters such as methyl acetate, ethyl acetate and others; halogenated hydrocarbons such as methylene chloride, chloroform, 1,2-dichloroethane and others; ethers such as diethyl ether, tetrahydrofuran, dioxane, ethylene glycol dimethyl ether, dithylene glycol dimethyl ether and others; ketones such as acetone, methyl ethyle ketone and others; aprotic polar solvents such as N,N-methylformamide, dimethyl sulfoxide, hexamethylphosphoryl triamide, N-methylpyrrolidone and others can be exemplified.

1 As to the ratio of the amount of the compound (26) to
the amount of the compound (27), generally at least an
equimolar quantity, preferably an equimolar quantity to
2 times the molar quantity of the latter may be used to
5 the former. The amount of the catalyst is not specifical-
ly restricted, and generally 0.01 to 10 times the molar
quantity, preferably 0.1 to 5 times the molar quantity
of the catalyst may be used to the compound (26). The
reaction is carried out generally at -20° to 200°C ,
10 preferably at -20° to 100°C , and is completed in 10
minutes to 20 hours.



Reaction process formula-14



[wherein R^5 is the same as defined above; and $\text{R}^{2'}$ is a lower alkyl group].

15 The reaction of a compound (26) with a known
compound (28) may be carried out under the same conditions
employed in the reaction of a compound (26) with a com-
pound (27) in the above-mentioned reaction process
formula-13.

1 Reaction process formula-15



[wherein R^5 , X^1 , M , A and R^6 are the same as defined above].

The halogenation reaction of a compound (4c)
 5 can be carried out in a suitable solvent, in the presence
 of a halogenating agent. As to the solvent used in the
 halogenation reaction, halogenated hydrocarbons such as
 chloroform, dichloromethane, carbon tetrachloride and
 others; ethers such as diethyl ether, tetrahydrofuran,
 10 dioxane and others; and acetic acid can be exemplified.
 As to the halogenating agent used in this halogenation
 reaction, halogen molecules such as bromine, chlorine and
 others; metal halides such as cupric bromide, cupric
 chloride, lithium chloride and others; thionyl chloride;
 15 N-halogenated succinimides such as N-chlorosuccinimide,
 N-bromosuccinimide and others can be exemplified.

The halogenating agent may generally be used
 in a large excess amount. The halogenation is carried
 out generally at 0° to 150°C , preferably at 0° to 120°C ,
 20 and is completed in 1 to 24 hours.

The reaction of a compound (29) with a compound


1 (30) can be carried out in a suitable solvent, in the
presence of a basic compound. As to the solvent used
in this reaction, ethers such as diethyl ether, dioxane,
tetrahydrofuran, ethylene glycol dimethyl ether, diethylene
5 glycol dimethyl ether and others; aromatic hydrocarbons
such as benzene, toluene, xylene and others; aliphatic
hydrocarbons such as n-hexane, heptane, cyclohexane and
others; aprotic polar solvents such as N,N-dimethyl-
formamide, dimethyl sulfoxide, hexamethylphosphoryl
10 triamide and others can be exemplified. As to the basic
compounds, inorganic basic compounds such as sodium
hydroxide, potassium hydroxide, sodium carbonate, potassium
carbonate, sodium hydrogen carbonate, potassium hydrogen
carbonate, metallic sodium, metallic potassium, sodium
15 amide, sodium hydride and others; alcoholates such as
sodium methylate, sodium ethylate and others; organic
basic compounds such as triethylamine, pyridine, N,N-
dimethylaniline and others can be exemplified.

The reaction is generally carried out at a room
20 temperature to 200°C, preferably at a room temperature to
150°C, and is completed in 1 to 24 hours. The ratio
of the amount of the compound (30) to the amount of the
compound (29) is generally at least an equimolar quantity,
preferably an equimolar quantity to 2 times the molar
25 quantity of the former is used to the latter.

Dihydropyridine derivatives represented by the
general formula (1) can also be prepared by the methods
as shown in the following reaction process formula-16

1 and -18.

Reaction process formula-16



[wherein R^1 , R^2 , R^3 , R^4 and X^1 are the same as defined above; E is a lower alkylene group; R^{14} is a 1,2,3,6-tetrahydropyridyl group which may have, as the substituent, a phenyl group which may have halogen atoms or lower alkyl groups as the substituents on the phenyl ring, group of the formula $\text{R}^6-(\text{D}')_{\text{O}}-\text{Y}-$ (wherein R^6 , D' , Y and O are the same as defined above)].


10 The reaction of a compound (31) with a compound (32) may be carried out in a suitable solvent, in the absence or presence of a basic compound. As to the solvent used in the reaction, halogenated hydrocarbons such as dichloromethane, dichloroethane, chloroform, carbon tetrachloride, and others; ethers such as diethyl ether, tetrahydrofuran, dioxane, ethylene glycol dimethyl ether and others; aliphatic hydrocarbons such as n-hexane, octane, cyclohexane and others; aromatic hydrocarbons such as benzene, toluene, xylene and others; aprotic polar

1 solvents such as acetone, acetonitrile, N,N-dimethyl-
formamide, dimethyl sulfoxide, hexamethylphosphoryl
triamide and others can be exemplified. As to the basic
compounds used in the reaction, inorganic basic compounds
5 such as sodium hydroxide, potassium hydroxide, sodium
carbonate, potassium carbonate, sodium hydrogen carbonate,
sodium hydride and others; amines such as triethylamine,
diisopropylethylamine, pyridine, quinoline and others
can be exemplified. Further, the reaction can be carried
10 out, if necessary, by adding an alkali metal iodide such
as potassium iodide, sodium iodide or the like, or
hexamethylphosphoryl triamide as the reaction accelerator.
The reaction is carried out at a room temperature to
200°C, preferably at a room temperature to 120°C, and is
15 completed in 1 to 24 hours.

The ratio of the amount of the compound (32) to
the amount of the compound (31) is at least an equimolar
quantity, preferably an equimolar quantity to 5 times the
molar quantity of the former to the latter.

20 Among the compounds represented by the general
formula (32), some of them are novel compounds and they
can be prepared by a method for example according to the
following reaction process formula-17.


1 Reaction process formula-17



[wherein R^8 , D' , $_o$ and X^1 are the same as defined above; and R^{15} is a lower alkanoyl group].

The reaction of a compound (33) with a compound (34) can be carried out under the same condition employed in the reaction of a compound (24) with a compound (25), and the hydrolysis followed by the reaction can be carried out under the same conditions employed in the hydrolysis of a compound (3g) in the above-mentioned reaction process formula-9.

Reaction process formula-18



1 [wherein R^1 , R^2 , R^3 , R^4 , E, X^1 , R^8 , D' and \underline{o} are the same as defined above; and X^2 is a halogen atom].

The reaction of a compound (31) with piperazine can be carried out under the same conditions employed

5 in the reaction of a compound (31) with a compound (32).

Further, the reaction of a compound (36) with a compound (34a) can be carried out under the same conditions employed in the reaction of a compound (24) with a compound (25).

10 Among the compounds represented by the general formula (1), those having basic groups can be converted into the corresponding salts by treating with pharmacologically acceptable acids. Examples of such acids including inorganic acids such as sulfuric acid, nitric
15 acid, hydrochloric acid, hydrobromic acid and others; as well as organic acids such as oxalic acid, maleic acid, fumaric acid, malic acid, citric acid, benzoic acid and others.

Compound of the present invention thus prepared can easily be isolated and purified by a method
20 usually employed in separation, such as precipitation, extraction, recrystallization, column chromatography, preparative thin layer chromatography and others.

Compound of the present invention represented
25 by the general formula (1) contains inevitably its optical isomers, as well as those in different crystal forms.

Compound of the present invention represented

1 by the general formula (1) can be administered, either
singly or together with conventional pharmacologically
acceptable carriers, to animals as well as to human
beings. No particular restriction is made to the
5 administration unit forms, thus compound of the present
invention represented by the general formula (1) can
be used in any desired administration unit form. Suitable
administration unit forms including peroral administration
unit forms such as tablets, granules and solutions;
10 and parenteral administration unit forms such as
injections.

Dosage of a compound represented by the general
formula (1) as the active ingredient to be administered
is not subjected to any particular restriction and can
15 be selected from a wide range. For the purpose of
attaining the desired pharmacological effects, it is
recommended to select a dosage from the range of 0.06
to 10 mg/kg of the body weight/day. It is suggested
also to contain 1 to 500 mg of the active ingredient in
20 each of the desired administration unit form.

In the present invention, the desired peroral
administration unit forms such as tablets, capsules
and solutions can be prepared by conventional methods.
For the purpose of shaping the administration unit form
25 into the form of tablets, a compound of the present
invention is mixed with pharmaceutically acceptable
excipients such as gelatin, starch, lactose, magnesium
stearate, talcum powder and gum arabic and others.

1 Capsules can be prepared by mixing a compound of the
present invention with an inert pharmaceutically accep-
table fillers or diluents and filling the mixture
obtained into rigid gelatin capsules or soft capsules.

5 Sirups or elixiers may be prepared by mixing a compound
of the present invention with a sweetening agent such
as sucrose; anticeptice such as methyl- or propyl-
parabens; colorants; seasoning agents and/or other
suitable additives. Parenteral preparations can also

10 be prepared by conventional methods, thus a compound
of the present invention is dissolved in a sterilized
liquid vehicle. As to the preferable vehicle, water
or saline can be used. Liquid preparations having desired
transparency, stability and parenteral use adaptability

15 can be prepared by dissolving approximately 1 to 500 mg
of the active ingredient in a solution of polyethylene
glycol having the molecular weight of 200 to 5,000, which
is soluble in both water and organic solvents. Desirably,
such liquid preparations may contain a lubricant such

20 as sodium carboxymethyl cellulose, methyl cellulose,
polyvinyl pyrrolidone and polyvinyl alcohol. Said
liquid preparations may also contain a bactericide and
fungicide such as benzyl alcohol, phenol and thimerosal,
and if necessary, an isotonic agent such as sucrose or

25 sodium chloride, a local anesthetic, stabilizer and buffer
solutions. Furthermore, additional ensurance of stability,
the parenteral compositions may be freezed after filling
and dehydrating steps by known lyophilization techniques.

- 1 The lyophilized powder of the parenteral composition can be made again into a normal use from just before the use.

Preparation of Tablets

1,000 Tablets for peroral use, each containing

- 5 5 mg of methyl 3-(4-hydroxyphenyl)-2-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate are prepared from the following formulation.

<u>Formulation</u>	<u>Amount (g)</u>
Methyl 3-(4-hydroxyphenyl)-2-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate	5
Lactose (Japanese Pharmacopoeia official drug grade)	50
Corn starch (Japanese Pharmacopoeia official drug grade)	25
Crystalline cellulose (Japanese Pharmacopoeia official drug grade)	25
Methyl cellulose (Japanese Pharmacopoeia official drug grade)	1.5
Magnesium stearate (Japanese Pharmacopoeia official drug grade)	1

- 10 Methyl 3-(4-hydroxyphenyl)-2-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate, lactose, corn starch and crystalline cellulose are mixed well, and the mixture is granulated with 5%-methyl cellulose aqueous solution, then the granules are passed through a 200 mesh sieve and then

- 1 dried carefully. The dried granules are passed through a 200 mesh sieve and mixed with magnesium stearate, then pressed into the form of tablets.

Preparation of capsules

- 5 1,000 Capsules of two-piece rigid gelatin capsules for peroral use, each containing 10 mg of methyl 3-(4-hydroxyphenyl)-2-propynyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate are prepared by using the following formulation.

<u>Formulation</u>	<u>Amount (g)</u>
Methyl 3-(4-hydroxyphenyl)-2-propynyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate	10
Lactose (Japanese Pharmacopoeia official drug grade)	80
Starch (Japanese Pharmacopoeia official drug grade)	30
Talcum powder (Japanese Pharmacopoeia official drug grade)	5
Magnesium stearate (Japanese Pharmacopoeia official drug grade)	1

- 10 The above-mentioned ingredients are finely ground, then mixed sufficiently to a uniform mixture and filled into gelatin capsules of a size having desired size for peroral administration.

1 Preparation of injection solution

A sterile aqueous solution suitable for parenteral use is prepared from the following formulation.

<u>Formulation</u>	<u>Amount (g)</u>
Methyl 3-(4-hydroxyphenyl)-2-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate	1
Polyethylene glycol (M. W. = 4,000) (Japanese Pharmacopoeia official drug grade)	0.9
Sodium chloride (Japanese Pharmacopoeia official drug grade)	0.9
Polyoxyethylene sorbitan monooleate (Japanese Pharmacopoeia official drug grade)	0.4
Sodium metabisulfite	0.1
Methyl p-hydroxybenzoate (Japanese Pharmacopoeia official drug grade)	0.18
Propyl p-hydroxybenzoate (Japanese Pharmacopoeia official drug grade)	0.02
Distilled water for injection	100 (ml)

The above-mentioned methyl p-hydroxybenzoate,
 5 propyl p-hydroxybenzoate, sodium metabisulfite and
 sodium chloride are dissolved in about a half volume of
 distilled water at 80°C under stirring condition. The
 solution obtained is cooled to 40°C, then methyl 3-(4-
 hydroxyphenyl)-2-propenyl 1,4-dihydro-2,6-dimethyl-4-
 10 (3-nitrophenyl)pyridine-3,5-dicarboxylate, polyethylene
 glycol and polyoxyethylene sorbitan monooleate are
 dissolved in the solution. Thus obtained solution is

1 further mixed with the distilled water for injection so
as to make it into the final volume, then sterilized by
sterile filtration with a suitable filter paper.

The present invention will be illustrated more
5 specifically by way of the following examples, in which
the preparations of the compounds to be used for the
starting materials will be shown in Reference Examples
and the preparations of the objective compounds will be
shown in Examples. The present invention, however will
10 not restricted to these examples.

Reference Example 1

To 20 g of p-hydroxybenzaldehyde and 32.5 g
of monoethyl malonate were added 6 ml of pyridine and
0.2 ml of piperidine, then the mixture thus obtained was
15 heated at 100 to 110°C for 10 hours under stirring.
The reaction mixture was then cooled, extracted with
chloroform, and the chloroform layer was washed with a
saturated aqueous solution of potassium hydrogen sulfite
and water in this order, the chloroform extract was
20 dried with magnesium sulfate. The solvent was removed
by evaporation and the residue thus obtained was crystal-
lized from isopropyl ether-n-hexane to yield 25.2 g of
ethyl 4-hydroxycinnamate. Light yellow indefinite form
crystals. Melting point: 70 - 71°C.

25 Reference Example 2

By using 20 g of 3-hydroxybenzaldehyde and

- 1 32.5 g of monoethyl malonate as the starting materials,
and by a method similar to that described in Reference
Example 1, under reaction conditions similar thereto,
there was prepared 25.5 g of ethyl 3-hydroxycinnamate.
5 Melting point: 65 - 68°C. (from isopropyl ether)

Reference Example 3

- By using 25.8 g of 4-hydroxy-3-chlorobenzaldehyde
and 32.5 g of monoethyl malonate as the starting materials,
and by a method similar to that described in Reference
10 Example 1, under reaction conditions similar thereto,
there was prepared 46 g of ethyl 4-hydroxy-3-chlorocin-
namate. Colorless prism-like crystals (from methylene
chloride). Melting point: 118 - 119°C.

Reference Example 4

- 15 To 30 ml of anhydrous ether solution containing
5 g of ethyl 4-hydroxycinnamate was added 7.1 ml of
dihdropyran and 50 mg of p-toluenesulfonic acid were
added, the mixture was stirred at a room temperature
for 2 hours, the reaction mixture was neutralized with
20 1%-sodium hydroxide solution, washed with water, and
dried with anhydrous sodium sulfate. The solvent was
removed by evaporation to yield 6.8 g of ethyl 4-(2-
tetrahydropyranyloxy)cinnamate. Colorless indefinite
form crystals. Melting point: 52 - 53°C.

1 Reference Example 5

50 Milliliters of anhydrous ether solution containing 6.8 g of ethyl 4-(2-tetrahydropyranyloxy)-cinnamate was added dropwise to an anhydrous ether solution containing 0.47 g of lithium aluminium hydride being cooled at -30°C . After the addition was finished, the reaction mixture was stirred for 1 hour at the same temperature, then the temperature of the reaction mixture was gradually elevated up to -10°C , then a saturated aqueous solution of sodium sulfate was gradually added to the reaction mixture, and the precipitates formed were removed by filtration. The filtrate was dried with anhydrous sodium sulfate, then concentrated to dryness, the residue thus obtained was treated by means of silica gel column chromatography (eluent: chloroform) to yield 3.2 g of 4-(2-tetrahydropyranyloxy)cinnamyl alcohol in the form of colorless oily substance. Refractive index: n_D^{22} 1.5520.

Reference Example 6

15 Grams of 4-(2-tetrahydropyranyloxy)cinnamyl alcohol and 5.2 g of sodium acetate were suspended in anhydrous methylene chloride, then to this suspension was added in one time 18 g of pyridium chlorochromate under ice-cooled condition. The reaction mixture was stirred for 1 hour at the same temperature, then the temperature was elevated to a room temperature, and the reaction mixture was stirred for additional 1 hour.

1 100 Milliliters of ether was added to the reaction
mixture, the whole mixture was filtered with Celite (a
trademark for diatomaceous product manufactured by and
sold from Johns-Manville Products Corp., Celite Division,
5 New York, N. Y., U. S. A.), the filtrate was concen-
trated and the residue thus obtained was treated by
silica gel column chromatography. Recrystallization
from ether to yield 3.5 g of 4-(2-tetrahydropyranyloxy)-
cinnamyl aldehyde in the form of colorless needle-like
10 crystals. Melting point: 65 - 67°C.

Reference Example 7

5.6 Grams of triethyl phosphonoacetate was
added dropwise under stirring condition at a room temper-
ature to a tetrahydrofuran solution containing 1.06 g
15 of 60%-sodium hydride, then the reaction mixture was
further stirred at 40°C for 1 hour. The reaction mixture
was cooled to a room temperature, then a tetrahydrofuran
solution containing 5.6 g of 4-(2-tetrahydropyranyloxy)-
cinnamyl aldehyde was added thereto, and stirred at a
20 room temperature for 2 hours, then the reaction mixture
was poured into 100 ml of water. The whole mixture was
extracted with ether, and the ether layer was washed
with water, and a saturated aqueous solution of sodium
chloride in this order, then dried with anhydrous sodium
25 sulfate. Recrystallization from isopropyl ether to
yield 3.8 g of ethyl 5-[4-(2-tetrahydropyranyloxy)phenyl]-
2(E),4(E)-pentadienoate in the form of colorless needle-like

1 crystals. Melting point: 66 - 67.5°.

Reference Example 8

To 30 ml of anhydrous benzene solution containing 3.6 g of ethyl 5-[4-(2-tetrahydropyranyloxy)phenyl]-
5 2(E),4(E)-pentadienoate was added dropwise 15 ml of diisobutyl aluminium hydride (25% by weight/by volume) under water-cooled condition and the reaction mixture was stirred at a room temperature for 2 hours. The reaction mixture was then poured into a saturated aqueous
10 solution of ammonium chloride and stirred at a room temperature for 2 hours. The insoluble matters were treated with Celite, and the insoluble matters were washed with ether. The organic layer was washed with water and dried with anhydrous sodium sulfate, then
15 concentrated to obtain the residue. The residue was recrystallized from chloroform-n-hexane to yield 2.8 g of 5-[4-(2-tetrahydropyranyloxy)phenyl]-2(E),4(E)-pentadienol in the form of colorless needle-like crystals. Melting point: 54 - 58°C.

20 Reference Example 9

25 Grams of p-hydroxyacetophenone, 50 ml of dihydropyrane and 0.25 g of p-toluenesulfonic acid were stirred at a room temperature for 2 hours in anhydrous ether. Then the reaction mixture was neutralized with
25 1N-sodium hydroxide and washed with water and a saturated sodium chloride aqueous solution in this order, then

1 dried with anhydrous sodium sulfate. The product was concentrated to yield 34 g of 4-(2-tetrahydropyranyloxy)-acetophenone. Colorless prism-like crystals. Melting point: 79 - 83°C.

5 Reference Example 10

45.8 Grams of triethyl phosphonoacetate and an anhydrous tetrahydrofuran solution containing 8.7 g of 60%-sodium hydride were stirred at 40°C for 1 hour, the reaction mixture was cooled and 30 g of 4-(2-tetra-
10 hydropyranyloxy)acetophenone was added to the reaction mixture. The whole reaction mixture was refluxed for 4 hours by heating, then the solvent was removed by evaporation, the residue thus obtained was extracted with ether and washed with water, than dried. The
15 extract was concentrated and the residue thus obtained was treated by means of silica gel column chromatography to yield 27.5 g of ethyl 3-methyl-p-(2-tetrahydropyranyloxy)cinnamate in the form of light yellow oily substance.

$^1\text{H-NMR}$ (90 MHz, CDCl_3) δ :

1.29 (3H, t, $J=6\text{Hz}$), 1.4 - 2.1 (6H, m),
2.49 (3H, d, $J=1\text{Hz}$), 3.3 - 3.9 (2H, m),
4.10 (2H, q, $J=6\text{Hz}$), 5.3 - 5.45 (1H, m),
6.03 (1H, d, $J=1\text{Hz}$), 6.9 - 7.4 (4H, m)

Reference Example 11

20 To a tetrahydrohydrofuran solution containing 27.5 g of ethyl 3-methyl-p-(2-tetrahydropyranyloxy)-

1 cinnamate was added dropwise 118 ml of diisobutyl
aluminium hydride (25% by weight/volume) at a room
temperature. After 2 hours, the reaction mixture was
poured into an ice-cooled ammonium chloride aqueous
5 solution, and the insoluble matters were removed by
filtration. The filtrate was washed with water then
dried with anhydrous sodium sulfate and concentrated.
The residue thus obtained was purified by means of silica
gel column chromatography to yield 12.7 g of 3-methyl-p-
10 (2-tetrahydropyranyloxy)cinnamyl alcohol in the form of
colorless oily substance.

¹H-NMR (90 MHz, CDCl₃) δ:

1.5 - 2.0 (6H, m), 2.0 (3H, s),
3.3 - 4.0 (2H, m), 4.1 - 4.3 (3H, m), 5.26 -
5.4 (1H, m), 5.81 (1H, t, J=6Hz), 6.8 - 7.5
(4H, m).

Reference Example 12

To a carbon disulfide solution containing 53 g
of aluminium chloride and 26.2 g of thioanisol was added
15 dropwise under an ice-cooled condition 24.0 ml of γ-
chlorobutyryl chloride. After 1 hour and 30 minutes,
the reaction mixture was poured into ice-water and the
insoluble matters were collected by filtration, dis-
solved in chloroform. The chloroform solution was
20 washed with water and dried with anhydrous sodium sulfate,
and concentrated to dryness. Recrystallization from
methanol to yield 36.2 g of γ-chloro-4-methylthiobutyro-

- 1 phenone in the form of light yellow prism-like crystals.
Melting point: 75 - 76°C.

Reference Example 13

- 35 Grams of γ -chloro-4-methylthiobutyrophenone,
5 34 ml of 1,8-diazabicyclo[5,4,0]undecene-7 (DBU) and
150 ml of acetonitrile were refluxed for 4 hours, then
500 ml of water was added to the reaction mixture, and
the whole mixture was extracted with ether. The ether
extract was washed with water, dried and the solvent
10 was removed by evaporation. The residue thus obtained
was recrystallized from methanol to yield 24.6 g of
cyclopropyl (4-methylthiophenyl)ketone as in the form
of light yellow prism-like crystals. Melting point:
76 - 76.5°C.

15 Reference Example 14

- To 200 ml of methanol solution containing 24 g
of cyclopropyl (4-methylthiophenyl)ketone was added
slowly 9.4 g of sodium borohydride under an ice-cooled
condition, the reaction mixture was stirred for 2 hours.
20 Then an adequate amount of acetone was added to the
reaction mixture and concentrated under a reduced pres-
sure. To the residue thus obtained was added chloroform,
and the chloroform solution was washed with water,
dried with anhydrous sodium sulfate, and the solvent was
25 removed by evaporation to yield 18 g of 1-(cyclopropyl,
hydroxymethyl)-4-methylthiobenzene as in the form of

1 colorless oily substance.

NMR (90 MHz, CDCl_3) δ :

0.2 - 0.7 (4H, m), 0.9 - 1.4 (1H, m), 1.4 - 2.0
(6H, m), 3.2 - 3.7 (2H, m), 5.17 - 5.3 (1H, m),
6.8 - 7.3 (8H, m).

Reference Example 15

To 10 ml of dioxane solution containing 5 g of
1-(cyclopropyl, hydroxymethyl)-4-methylthiobenzene was
5 added dropwise 6 ml of 47% of hydrobromic acid under an
ice-cooled condition, then the mixture was stirred for
30 minutes, and the reaction mixture was concentrated
under a reduced pressure. To the residue thus obtained
was added water, then extracted with ether, the extract
10 was washed with water and dried with anhydrous sodium
sulfate, and concentrated to dryness. The residue was
recrystallized from methanol to yield 2.2 g of 4-(4'-
methylthiophenyl)-3(E)-butenylbromide in the form of
colorless flake-like crystals. Melting point: 54 - 56°C.

15 Reference Example 16

10 Grams of p-iodophenol, 8 ml of dihydropyran
and a catalytic amount of p-toluenesulfonic acid was
dissolved in 30 ml of anhydrous ether, all of these were
mixed together and stirred at a room temperature for 2
20 hours. The reaction mixture was washed with water,
dried then the solvent was removed by evaporation to
yield 12.4 g of 4-(2-tetrahydropyranyloxy)-1-iodobenzene

1 in the form of yellow oily substance. Boiling point:
84 - 87°C (at 25 mm Hg).

Reference Example 17

6.25 Grams of triethyl phosphonocrotonate was
5 added dropwise to a tetrahydrofuran solution containing
1.06 g of 60%-sodium hydride at a room temperature, the
reaction mixture was stirred for 1 hour at 40°C. The
temperature of the reaction mixture was cooled to a room
temperature, then a tetrahydrofuran solution containing
10 5.0 g of p-(2-tetrahydropyranyloxy)benzaldehyde was
added to the reaction mixture, and stirred at a room
temperature for 2 hours, then poured into 100 ml of
water. The whole mixture was extracted with ether,
the ether extract was washed with water and a saturated
15 sodium chloride aqueous solution in this order, then
dried with anhydrous sodium sulfate. The solvent was
removed by evaporation, the residue thus obtained was
purified by means of silica gel column chromatography
(eluent: hexane-chloroform), next recrystallized from
20 isopropyl ether to yield 4.01 g of ethyl 5-[4-(2-tetra-
hydropyranyloxyphenyl)]-2(E),4(E)-pentadienoate. Colorless
needle-like crystals. Melting point: 66 - 67.5°C.

Reference Example 18

20 Grams of p-tetrahydropyranyloxyiodobenzene
25 and 70 ml of anhydrous pyridine containing 11.3 g of
copper (I) 3-acetyloxy-1-propyn-1-ide were refluxed for

1 6 hours under an atmosphere of argon gas. After the
reaction was finished the reaction mixture was poured
into water and the whole mixture was extracted with
chloroform. The chloroform layer was washed with water,
5 dried and the solvent was removed by evaporation. The
residue thus obtained was purified by means of a silica
gel column chromatography (eluent: chloroform: n-hexane =
1:1) to yield 8 g of 4-[4-(2-tetrahydropyranyloxy)phenyl]-
3-butynyl acetate in the form of colorless oily substance.

¹H-NMR (60 MHz, CDCl₃) δ:

6.98 (2H, d, J=8Hz), 6.63 (2H, d, J=8Hz),
6.15 (1H, m), 4.05 (2H, t, J=6Hz), 3.3 - 3.7
(2H, m), 2.58 (2H, t, J=6Hz), 1.97 (3H, s),
1.5 - 1.9 (6H, m).

10 Reference Example 19

To an anhydrous tetrahydrofuran solution contain-
ing 2.4 g of 4-[4-(2-tetrahydropyranyloxy)phenyl]-
3-butynylacetate was added 1 g of lithium aluminium
hydride and the mixture was refluxed for 12 hours. After
15 the reaction was finished, a saturated sodium sulfate
aqueous solution was slowly added to the reaction mix-
ture, and the precipitates formed were removed by
filtration, then the filtrate was dried with anhydrous
sodium sulfate, and concentrated to dryness. The residue
20 thus obtained was purified by means of a silica gel
column chromatography to yield 2 g of 4-[(2-tetrahydro-
pyranyloxy)phenyl]-3(E)-butenylalcohol as in the form

1 of colorless oily substance.

¹H-NMR (90 MHz, CDCl₃) δ:

1.5 - 2.1 (6H, m), 2.43 (2H, q, J=6Hz),
3.4 - 4.0 (5H, m), 5.37 (1H, m), 6.03 (1H, d, t,
J=16Hz, 6Hz), 6.40 (1H, d, J=16 Hz), 6.97 (2H,
d, J=9Hz), 7.25 (2H, d, J=9Hz).

Reference Example 20

15 Grams of 1-iodo-4-(1-ethoxyethoxy)benzene,
6 g of propargyl acetate, 0.26 g of triphenylphosphine,
5 0.09 g of palladium chloride and 20 ml of diethylamine
solution of cuprous iodide were stirred at 40 - 50°C for
1 hour. After the reaction was finished, the reaction
mixture was poured into water, and extracted with diethyl
ether. The ether extract was washed with water, dried
10 and the solvent was removed by evaporation, then the
residue thus obtained was purified by means of a silica
gel column chromatography (eluent: dichloromethane:
n-hexane = 1:1 and dichloromethane) to yield 10.1 g of
3-[4-(1-ethoxyethoxy)phenyl]propargyl acetate.

N. M. R. (CDCl₃) δ:

1.17 (3H, t, J=7Hz), 1.47 (3H, d, J=5Hz),
2.09 (3H, s), 3.3 - 3.9 (2H, m), 4.87 (2H,
s), 5.37 (1H, q, J=5Hz), 6.92 (2H, d, J=9Hz),
7.37 (2H, q, J=9Hz).

15 Reference Example 21

To 50 ml of methanol solution containing 10 g

1 of 3-[4-(1-ethoxyethoxy)phenyl]propargyl acetate, were
added 1.5 g of 5%-Pd-BaSO₄ and 10 drops of quinoline,
and the mixture was catalytically reduced. After the
reaction was finished, the reaction mixture was filtered,
5 and the filtrate was allowed to evaporation, the residue
thus obtained was purified by means of a silica gel
column chromatography (eluent: dichloromethane and
dichloromethane:methanol = 100:1) to obtain 3 g of
4(Z)-(1-ethoxyethoxy)cinnamyl acetate and 4 g of 4(Z)-
10 hydroxycinnamyl acetate. Thus obtained 4 g of 4(Z)-
hydroxycinnamyl acetate was dissolved in 20 ml of
anhydrous ether, to this solution was added 50 mg of p-
toluenesulfonic acid and 10 ml of ethyl vinyl ether, and
the whole mixture was refluxed for 3 hours. The reaction
15 mixture was then washed with 5%-sodium hydroxide aqueous
solution and water in this order, and dried. The solvent
was removed by evaporation to yield 4.8 g of 4(Z)-(1-
ethoxyethoxy)cinnamyl acetate.

N. M. R. (CDCl₃) δ:

1.21 (3H, t, J=7Hz), 1.50 (3H, d, J=5Hz),
2.09 (3H, s), 3.24 - 3.64 (1H, m), 3.70 - 3.90
(1H, m), 4.84 (2H, dd, J=6.5Hz, 1.5Hz), 5.40
(1H, q, J=5Hz), 5.73 (1H, dt, J=11.5Hz,
6.5Hz), 6.60 (1H, dm, J=11.5Hz), 6.98 (2H, d,
J=9Hz), 7.15 (2H, d, J=9Hz).

Reference Example 22

20 7.8 Grams of 4(Z)-(1-ethoxyethoxy)cinnamyl

1 acetate and 50 ml of methanol solution containing 5 g of
potassium carbonate were stirred at a room temperature
for 2 hours. The solvent was removed by evaporation,
and to the residue thus obtained was added water and
5 extracted with ether. The ether extract was washed with
water, dried and the solvent removed by evaporation to
yield 4(Z)-(1-ethoxyethoxy)cinnamyl alcohol.

N. M. R. (90 MHz, CDCl_3) δ :

1.19 (3H, t, $J=7\text{Hz}$), 1.48 (3H, d, $J=5\text{Hz}$),
3.4 - 3.9 (2H, m), 4.39 (2H, dd, $J=6\text{Hz}$, $J=2\text{Hz}$),
5.36 (1H, q, $J=5\text{Hz}$), 5.75 (1H, dt, $J=11\text{Hz}$,
 $J=6\text{Hz}$), 6.45 (1H, dm, $J=11\text{Hz}$), 6.95, 7.10 (4H,
AB-q, $J=9\text{Hz}$).

Reference Example 23

8 Milliliters of 85% sodium hydroxide aqueous
10 solution mixed with 46 g of ethylene glycol was heated
at 40 - 50°C, and 15 g of cinnamyl bromide was added
dropwise thereto. After the addition was finished, the
reaction mixture was heated at 100 - 110°C for 2 hours.
The reaction mixture was allowed to cool, and poured
15 into 50 ml of water, then extracted with ether. The
ether extract was washed with water and a saturated
aqueous solution of sodium chloride in this order, and
dried with anhydrous sodium sulfate. The solvent was
removed by evaporation, the residue thus obtained was
20 purified by means of a silica gel column chromatography
(eluent: chloroform) to yield 5.5 g of 2-[3-phenyl-

1 2(E)-propenyloxy]ethanol.

N. M. R. CDCl_3 (90 MHz) δ :

2.20 (1H, t, $J=6\text{Hz}$), 3.50 - 3.85 (4H, m),
4.18 (2H, d, $J=6\text{Hz}$), 6.23 (1H, dt, $J=6\text{Hz}$,
 $J=16\text{Hz}$), 6.57 (1H, d, $J=16\text{Hz}$), 7.20 - 7.50
(5H, m).

Reference Example 24

5.33 Grams of 60% sodium hydride was suspended
in 25 ml of dimethylformamide, and 9.3 ml of ethylene
5 glycol was added dropwise thereto. After the addition
was finished the reaction mixture was stirred at 45°C
for 2 hours. The reaction mixture was ice-cooled, then
6.5 g of 3-phenylpropargyl bromide was added dropwise to
the reaction mixture and stirred at the same temperature
10 for 1 hour. After the reaction was finished, the reaction
mixture was poured into 200 ml of water, and the whole
mixture was extracted with ether. The ether extract
was washed with water and dried with anhydrous magnesium
sulfate, and the solvent was removed by evaporation. The
15 residue thus obtained was purified by means of a silica
gel column chromatography (eluent: chloroform:n-hexane =
3:1) to yield 2.4 g of 2-(3-phenyl-2-propionyloxy)ethanol.
Refractive index: n_D^{24} 1.5590.

Reference Example 25

20 6.5 Grams of propargyl bromide, 0.32 g of tetra-
butylammonium bromide and 2.7 g of sodium hydroxide were

1 dissolved in 20 ml of methylene chloride and 8 ml of
water, and this solution was refluxed for 2 hours. After
the reaction mixture was cooled, the organic layer was
collected by separation, washed with water, dried with
5 anhydrous magnesium sulfate. The solvent was removed by
evaporation, the residue thus obtained was purified by
means of a silica gel column chromatography (eluent:
chloroform:n-hexane = 2:1) to yield 4.0 g of 2-(3-
phenyl-2-propionylthio)ethanol. Refractive index:
10 n_D^{22} 1.6078.

Reference Example 26

4.3 Grams of 2-mercaptoethanol, 10 ml of 40%
sodium hydroxide aqueous solution, 0.2 g of tetrabutyl-
ammonium bromide and 40 ml of methylene chloride were
15 heated at 40 - 45°C, then 20 ml of methylene chloride
solution containing 10 g of cinnamyl bromide was added
dropwise thereto. The reaction mixture was stirred
vigorously at the same temperature for 4 - 5 hours. The
organic layer was collected by separation and washed with
20 water, dried with anhydrous sodium sulfate. The solvent
was removed by evaporation, the residue thus obtained
was distilled under a reduced pressure to yield 8.6 g
of 2-(3-phenyl-2(E)-propionylthio)ethanol. Boiling
point: 150 - 153°C (1 mm Hg).

N. M. R. $CDCl_3$ (90 MHz) δ :

2.16 (1H, t, J=6Hz), 2.70 (2H, t, J=6Hz),

3.30 (2H, d, J=6Hz), 3.70 (2H, q, J=6Hz),

6.14 (1H, dt, J=6Hz, J=16Hz), 6.40 (1H, d, J=16Hz), 7.20 - 7.60 (5H, m).

1 Reference Example 27

12.5 Grams of cinnamyl chloride and 21 g of 2-methylaminoethanol were dissolved in 100 ml of ethanol, and the solution was refluxed for 5 hours. The solvent was removed by evaporation, to the residue thus obtained was added water and washed with n-hexane. To the water layer was added 3N-sodium hydroxide aqueous solution to adjust its pH to 10 - 11, and extracted with ether. The ether extract was washed with water and dried with anhydrous sodium sulfate, and the solvent was removed by evaporation. The residue was purified by means of a silica gel column chromatography (eluent: chloroform: methanol = 25:1) to yield 15 g of 2-[N-methyl-N-(3-phenyl-2(E)-propenyl)amino]ethanol.

N. M. R. CDCl_3 (90 MHz) δ :

2.30 (3H, s), 2.55 (2H, t, J=7Hz), 2.90 (1H, s), 3.20 (2H, d, J=7Hz), 3.60 (2H, t, J=7Hz), 6.20 (1H, dt, J=7Hz, J=16Hz), 6.48 (1H, d, J=16Hz), 7.25 - 7.50 (5H, m).

15 Reference Example 28

By a method similar to that described in Reference Example 27, by using a suitable starting material, the following compound was prepared.

2-[N-Methyl-N-(3-phenyl-2-propynyl)amino]ethanol

Boiling point: 124 - 126°C (1mm Hg)

Refractive index: n_D^{24} 1.5589

1 Reference Example 29

5 Grams of cinnamyl bromide, 5 g of anhydrous piperazine and 50 ml of acetone solution containing 8.4 g of anhydrous potassium carbonate were refluxed for 5 6 hours. To the reaction mixture was added water and the whole mixture was extracted with chloroform. The chloroform layer was washed with water, dried and the solvent was removed by evaporation. 2.2 Grams of 1-acetyl-4-cinnamylpiperazine thus obtained was dissolved 10 in 80% of methanol, and 2 g of potassium hydroxide was added thereto, then the mixture was refluxed for 8 hours. To the reaction mixture was added water, and the whole mixture was extracted with chloroform, washed with water, dried, and chloroform was removed by evaporation. The 15 residue thus obtained was purified by means of a silica gel column chromatography (eluent: chloroform and chloroform:methanol = 30:1) to yield 1.2 g of N-cinnamylpiperazine.

N. M. R. $CDCl_3$ δ :

2.34 (1H, s), 2.46 (4H, m, $J=5\text{Hz}$), 2.92 (4H, m, $J=5\text{Hz}$), 3.13 (2H, d, $J=6\text{Hz}$), 6.23 (1H, dt, $J=15\text{Hz}$, 6Hz), 6.82 (1H, d, $J=15\text{Hz}$), 7.2 - 7.5 (5H, m).

1 Reference Example 30

To 150 ml of anhydrous carbon tetrachloride solution containing 116 g of methyl acetoacetate was added dropwise 51 ml of bromine at below 5°C. After the addition was finished the reaction mixture was stirred overnight, then the reaction mixture was poured into ice-water, washed with a diluted sodium carbonate aqueous solution five times, then with a saturated sodium chloride aqueous solution, and dried with anhydrous calcium chloride. The solvent was removed by evaporation to yield 170 g of methyl 4-bromoacetoacetate.

Reference Example 31

1.6 Grams of 60% sodium hydride was suspended in 80 ml of anhydrous dimethoxyethane, then 40 ml of dimethoxyethane solution containing 8 g of methyl 4-bromoacetoacetate was added dropwise thereto at -40 to -50°C. At the same temperature, lithium N-methylcinnamylamide prepared from 20 ml of dimethoxyethane solution containing 5.9 g of N-methylcinnamylamine and 40 ml of 1N-n-butyl lithium was added dropwise. After the addition was finished, the reaction mixture was stirred at 65 - 70°C for 15 hours. The reaction mixture was poured into an ice-water and the whole mixture was extracted with diethyl ether. The extract was washed with water, dried and the solvent was removed by evaporation, the residue thus obtained was purified by means a silica gel column chromatography (eluent: dichloromethane and dichloro-

1 methane:methanol = 50:1) to yield 1.8 g of 4-(N-methyl-
E-cinnamylamino)acetoacetate.

N. M. R. CDCl_3 δ :

2.32 (3H, s), 3.22 (2H, d, $J=6\text{Hz}$), 3.31 (2H, s),
3.51 (2H, 2), 3.68 (3H, s), 6.18 (1H, dt,
 $J=16\text{Hz}$, 6Hz), 6.50 (1H, d, $J=16\text{Hz}$), 7.0 - 7.5
(5H, m).

Reference Example 32

4 Grams of 60% sodium hydride was suspended in
5 150 ml of anhydrous dimethoxyethane and the suspension
was cooled to -30°C . Then, 19.4 g of methyl 4-bromo-
acetoacetate was added dropwise to the suspension and
stirred for 1 hour. Further, at the same temperature, an
anhydrous dimethoxyethane solution of sodium cinnamy
10 alcoholate prepared from 13.4 g of cinnamyl alcohol, 4 g
of 60% of sodium hydride and 75 ml of dimethoxyethane
was added dropwise thereto, and the reaction mixture was
refluxed for 15 hours. After the reaction mixture was
cooled to a room temperature, the reaction mixture was
15 poured into 1N-hydrochloric acid being ice-cooled, and
the whole mixture was extracted with ether. The ether
extract was washed with water and dried with anhydrous
sodium sulfate, and the solvent was removed by evaporation.
The residue was purified by means of a silica gel column
20 chromatography (eluent: methanol:dichloromethane = 1:20)
to yield 6 g of methyl 4(E)-cinnamyloxyacetoacetate
Refractive index: n_D^{20} 1.5402

1 Reference Example 33

By a method similar to that described in Example 32, by using a suitable starting material, there was prepared the following compounds.

Methyl 4-[4(E)-(1-ethoxyethoxy)cinnamyloxy]-
acetoacetate

Refractive index: $n_D^{20} = 1.5232$

5 Reference Example 34

A mixture of 9 g of 4(E)-(1-ethoxyethoxy)-
cinnamyl alcohol and 0.5 ml of triethylamine was heated
at 70 - 80°C, and 6 g of diketene was slowly added dropwise
thereto. After the addition was finished, the reaction
10 mixture was heated at 110 - 120°C for 30 minutes. After
the reaction was completed, the solvent was removed by
evaporation under a reduced pressure, and the residue
thus obtained was purified by means of a silica gel
column chromatography (eluent: diethyl ether: n-hexane =
15 1:3) to yield 8.8 g of 4(E)-(1-ethoxyethoxy)cinnamyl
acetoacetate. Refractive index: $n_D^{24} = 1.5305$

N. M. R. $CDCl_3$ (90 MHz) δ :

1.17 (3H, t, J=7Hz), 1.48 (3H, d, J=6Hz),
2.25 (3H, s), 3.48 (2H, s), 3.40 - 3.85 (2H, m),
4.75 (2H, d, J=6Hz), 5.36 (1H, q, J=6Hz),
6.18 (1H, dt, J=6Hz, J=16Hz), 6.94 (2H, d,
J=9Hz), 7.28 (2H, d, J=9Hz).

1 Reference Example 35

To 400 ml of acetonitrile solution containing 30 g of 2-iodoethyl methyl 2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate was added 26.6 g of piperazine, and the mixture was stirred at a room temperature overnight. After the reaction was finished, the insoluble matters were removed by filtration, then chloroform was added to the filtrate, the chloroform layer was washed with a saturated sodium hydrogen carbonate aqueous solution and water in this order, and dried with anhydrous sodium sulfate. The reaction mixture was concentrated by evaporating the solvent, and the residue thus obtained was purified by means of a silica gel column chromatography (eluent: dichloromethane:methanol:triethylamine = 100:10:1) to yield 21 g of methyl 5-[2-(1-piperazinyl)ethyl] 2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate as in the form of light yellow indefinite form crystals.

N. M. R. CDCl_3 (200 MHz) δ :

1.892 (1H, bs), 2.366 (3H, s), 2.379 (3H, s),
2.40 - 2.50 (4H, m), 2.52 - 2.60 (2H, m),
2.80 - 2.90 (4H, m), 3.651 (3H, s), 4.08 -
4.28 (2H, m), 5.109 (1H, s), 5.947 (1H, bs),
7.3 - 8.2 (4H, m).

Reference Example 36

20 18.6 Grams of 5-acetyl-2,2-dimethyl-1,3-dioxane-4,6-dione and 50 ml tetrahydrofuran solution containing

- 1 22.2 g of 4-(1-ethoxyethoxy)cinnamyl alcohol were refluxed for 6 hours. After the reaction was completed, the solvent was removed by evaporation, and the residue thus obtained was purified by means of a silica gel
- 5 column chromatography (eluent: ethyl acetate:n-hexane = 1:4) to yield 5.8 g of 4-[(1-ethoxy)ethoxy]cinnamyl acetoacetate as in the form of an oily substance.

N. M. R. (CDCl_3) δ :

1.15 (3H, t, $J=7\text{Hz}$), 1.44 (3H, d, $J=5.4\text{Hz}$),
2.20 (3H, s), 3.31 - 3.95 (2H, m), 3.45 (2H,
s), 4.71 (2H, d, $J=6\text{Hz}$), 5.35 (1H, q, $J=5.2\text{Hz}$),
6.10 (1H, dt, $J=6.1\text{Hz}$, 15.3Hz), 6.59 (1H, d,
 $J=15.3\text{Hz}$), 6.91 (2H, d, $J=9\text{Hz}$), 7.29 (2H, d,
 $J=9\text{Hz}$).

IR (film): ν : 1650, 1730^{-1} .

Example 1

- 5.4 Grams of 4-(2-tetrahydropyranyloxy)cinnamyl
- 10 alcohol, 11.5 g of 1,4-dihydro-2,6-dimethyl-5-methoxy-carbonyl-4-(3-nitrophenyl)pyridine-3-carboxylic acid and 7.1 g of DCC were dissolved in 30 ml of pyridine, and the reaction mixture was stirred at a room temperature overnight. 200 Milliliters of water was added to the
- 15 reaction mixture and extracted with ethyl acetate. The extract was washed with water, a saturated potassium hydrogen sulfate aqueous solution, water and a saturated sodium chloride aqueous solution in this order, then dried and the solvent was removed by evaporation. The

1 residue thus obtained was purified by means a silica gel
column chromatography (eluent: chloroform) to yield
3.4 g of methyl 3-(4-tetrahydropyranyloxyphenyl)-2(E)-
propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-
5 pyridine-3,5-dicarboxylate.

N. M. R. δ (CDCl_3):

1.4 - 2.0 (6H, m), 2.26 (6H, m), 3.53 (3H, s),
3.4 - 4.0 (2H, m), 4.58 (2H, d, $J=6\text{Hz}$), 5.03
(1H, s), 5.30 (1H, t, $J=3\text{Hz}$), 5.97 (1H, dt,
 $J_a=6\text{Hz}$, $J_b=16\text{Hz}$), 6.30 (1H, bs), 6.37 (1H, d,
 $J=16\text{Hz}$), 6.88 (2H, d, $J=9\text{Hz}$), 7.14 (2H, d,
 $J=9\text{Hz}$), 7.20 (1H, t, $J=6\text{Hz}$), 7.50 (1H, dt,
 $J_a=2\text{Hz}$, $J_b=6\text{Hz}$), 7.82 (1H, dt, $J_a=2\text{Hz}$, $J_b=t\text{Hz}$),
8.00 (1H, t, $J=2\text{Hz}$).

Example 2

3.4 Grams of methyl 3-(4-tetrahydropyranyloxy-
phenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-
nitrophenyl)pyridine-3,5-dicarboxylate was dissolved in
10 50 ml of methanol, then 0.2 g of *p*-toluenesulfonic acid
was added to the solution and the whole mixture was
stirred at a room temperature for 4 hours. The reaction
mixture was neutralized by adding sodium hydrogen
carbonate, and methanol was removed by evaporation. The
15 residue thus obtained was purified by means of a silica
gel column chromatography (eluent: chloroform), recrystal-
lized from benzene-ether to yield 2 g of methyl 3-(4-
hydroxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-

- 1 (3-nitrophenyl)pyridine-3,5-dicarboxylate as in the form
of light yellow powdery substance. Melting point: 137.5 -
139°C.

Example 3

- 5 To 20 ml of an anhydrous methylene chloride
solution containing 1.6 g of tetramethylurea was added
1.9 g of oxalyl chloride at a room temperature and the
mixture was refluxed for 2 hours, then 4.3 g of 1,4-
dihydro-2,6-dimethyl-5-methoxycarbonyl-4-(3-nitrophenyl)-
10 pyridine-3-carboxylic acid was added thereto, under an
ice-cooled and stirred condition while the reaction
mixture was as in a suspended state, 1.7 g of 3-p-(2-
tetrahydropyranyloxy)phenylpropargyl alcohol in 10 ml of
anhydrous methylene chloride solution was added to the
15 suspension and the whole mixture was stirred at a room
temperature for 3 hours. Under an ice-cooled condition,
the reaction mixture was poured in 1N-hydrochloric acid,
and the organic layer was washed with water in three
times and dried with anhydrous sodium sulfate, then
20 solvent was removed by evaporation. The residue thus
obtained was purified by means of a silica gel column
chromatography (eluent: chloroform) to yield 2.8 g of
methyl 3-[4-(2-tetrahydropyranyloxyphenyl)]-2-propynyl
1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-
25 dicarboxylate in the form of light yellow indefinite
form crystals.

1 Example 4

2.7 Grams of methyl 3-[4-(2-tetrahydropyranyloxy-phenyl)]-2-propynyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate was dissolved in
5 30 ml of methanol, then 10 mg of p-toluenesulfinic acid was added and the mixture was stirred for 1 hour. The reaction mixture was neutralized by adding a saturated sodium hydrogen carbonate aqueous solution, further 100 ml of water was added then the mixture was extracted with
10 chloroform. The chloroform layer was washed with water in three times, then dried with anhydrous sodium sulfate, the solvent was removed by evaporation. The residue thus obtained was purified by means of a silica gel column chromatography (eluent: ethyl acetate:n-hexane = 4:1),
15 next recrystallized from tetrahydrofuran-n-hexane to yield 1.2 g of methyl 3-(4-hydroxyphenyl)-2-propynyl-1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate in the form of yellow prism-like crystals. Melting point: 173 - 176°C.

20 Example 5

To a mixture consisting of 1.9 of 1,4-dihydro-2,6-dimethyl-5-methoxycarbonyl-4-(3-nitrophenyl)pyridine-3-carboxylic acid, 0.78 ml of 30% sodium hydroxide aqueous solution and 15 ml of HMPA were added dropwise 1.5 g of
25 4-(4-methylthiophenyl)-3(E)-butenyl bromide. The mixture was stirred at a room temperature for 8 hours, then at 40 - 45°C for 5 hours, next the reaction mixture was

1 poured into an ice-water and extracted with chloroform.
The chloroform layer was washed with water, then dried
with anhydrous sodium sulfate, and the solvent was
removed by evaporation. The residue thus obtained was
5 purified by means of a silica gel column chromatography
(eluent: ethyl acetate:n-hexane = 1:2), recrystallization
from ethyl acetate-n-hexane to yield 1.0 g of methyl
4-(4-methylthiophenyl)-3(E)-butenyl 1,4-dihydro-2,6-
dimethyl-4-(3-nitrophenyl)-pyridine-3,5-dicarboxylate in
10 the form of light yellow powdery substance. Melting
point: 165 - 170°C.

Example 6

3.36 Grams of 1-[2(E)-acetoacetocymethylvinyl]-
4-(2-tetrahydropyranyloxy)benzene, 1.5 g of 3-nitro-
15 benzaldehyde and 1.2 g of methyl 3-aminocrotonate
were added to 20 ml of isopropanol, and the mixture was
refluxed for 8 hours. The reaction mixture was
concentrated and the residue was purified by means of a
silica gel column chromatography (eluent: chloroform)
20 to yield 1.2 g of methyl 3-(4-tetrahydropyranyloxyphenyl)-
2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-
pyridine-3,5-dicarboxylate.

N. M. R. δ (CDCl₃):

1.4 - 2.0 (6H, m), 2.26 (6H, m), 3.53 (3H, s),
3.4 - 4.0 (2H, m), 4.58 (2H, d, J=6Hz), 5.03
(1H, s), 5.30 (1H, t, J=3Hz), 5.97 (1H, dt,
Ja=6Hz, Jb=16H), 6.30 (1H, bs), 6.37 (1H, d,


J=16Hz), 6.88 (2H, d, J=9Hz), 7.14 (2H, d, J=9Hz), 7.20 (1H, t, J=6Hz), 7.50 (1H, dt, Ja=2Hz, Jb=6Hz), 7.82 (1H, dt, Ja=2Hz, Jb=6Hz), 8.00 (1H, t, J=2Hz).

1 Examples 7 - 30

By using a suitable starting material, and by methods similar to those described in Examples 1 and 6, there were prepared compounds as shown in Table 1 as

5 follows.

Table 1











Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
7	CH ₃	CH ₃		CH ₃		137.5 - 139 (Ether-benzene)	yellow powdery crystals IR ¹)	-
8	CH ₃	CH ₃		CH ₃		60 - 63 (Ether-n-hexane)	Light yellow powdery crystals	-
9	CH ₃	CH ₃		CH ₃		75 - 105 (Ether-n-hexane)	Light yellow indefinite form crystals	-

(to be continued)

Table 1









(Continued)

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
10	CH ₃	CH ₃		CH ₃		181 - 182 (Methanol)	yellow powdery crystals	-
11	CH ₃	CH ₃		CH ₃		173 - 176 (Tetrahydrofuran- n-hexane)	yellow prism-like crystals	-
12	CH ₃	CH ₃		CH ₃		170 - 171 (Methanol)	Light yellow powdery crystals	-
13	CH ₃	CH ₃		CH ₃		90 - 95 (Tetrahydrofuran- n-hexane)	Light yellow indefinite form crystals	-

(to be continued)

Table 1

(Continued)









Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
14	CH ₃	CH ₃		CH ₃		164 - 165 (Methanol-ether)	Light yellow powdery crystals	-
15	CH ₃	CH ₃		CH ₃		146 - 153 (Isopropanol)	Yellow powdery crystals	-
16	CH ₃	CH ₃		CH ₃		165 - 170 (Ethyl acetate -n-hexane)	Light yellow indefinite form crystals	-
17	CH ₃	CH ₃		CH ₃		145 - 150 (Ethyl acetate -n-hexane)	Yellow indefinite form crystals	-

(to be continued)

0145434

Table 1

(Continued)









Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
18	CH ₃	CH ₃		CH ₃		68 - 70 (Ether-n-hexane)	Light yellow powdery crystals	-
19	CH ₃	CH ₃		CH ₃		NMR ¹⁾	Light yellow indefinite form crystals	-
20	CH ₃	CH ₃		CH ₃		NMR ²⁾	Light yellow indefinite form crystals	-
21	CH ₃	CH ₃		CH ₃		NMR ³⁾	Light yellow indefinite form crystals	-

(to be continued)

(*) OTHP: 2-tetrahydropyranyloxy group

Table 1









(Continued)

Example No.	¹ R	² R	³ R	⁴ R	⁵ R	Melting point (°C) (Recrystallization Solvent)	Crystal form	Salt
22	CH ₃	CH ₃		CH ₃	 OTHP (*)	NMR ⁴⁾	Light yellow indefinite form crystals	-
23	CH ₃	CH ₃		CH ₃	 OTHP (*)	NMR ⁵⁾	Light yellow indefinite form crystals	-
24	CH ₃	CH ₃		CH ₃	 OTHP (*)	NMR ⁶⁾	Light yellow indefinite form crystals	-
25	CH ₃	CH ₃		CH ₃	 OTHP (*)	NMR ⁷⁾	Yellow indefinite form crystals	-

(*) OTHP: 2-tetrahydropyranyloxy group

(to be continued)









Table 1

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
26	CH ₃	CH ₃		CH ₃	 (*)	NMR ⁸⁾	Light yellow indefinite form crystals	-
27	CH ₃	CH ₃		CH ₃	 (*)	NMR ⁹⁾	Light yellow indefinite form crystals	-
28	CH ₃	CH ₃		CH ₃	 NMR ¹⁰⁾	NMR ¹⁰⁾	Light yellow indefinite form crystals	-
29	CH ₃	CH ₃		CH ₃		120.5 - 121.5 (Tetrahydrofuran-n-hexane)	Yellow powdery crystals	-

(*) OTHP: 2-tetrahydropyranyloxy group (to be continued)

Table 1

(Continued)









Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
30	CH ₃	CH ₃		CH ₃		122 - 124 (Tetrahydrofuran- n-hexane)	Light yellow powdery crystals	-
31	CH ₃	CH ₃		CH ₃		92 - 95 (Chloroform- n-hexane)	Light yellow powdery crystals	-
32	CH ₃	CH ₃		CH ₃		NMR ¹²⁾	Yellow oily substance	-
33	CH ₃	CH ₃		CH ₃		170 - 172 (Ether- n-hexane)	Light yellow prism-like crystals	-

(*) OTHP: 2-tetrahydropyranyloxy group

(to be continued)

Table 1










(Continued)

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
34	CH ₃	CH ₃		CH ₃		181 - 182 (Acetone)	Light yellow prism-like crystals	-
35	CH ₃	CH ₃		CH ₃		168 - 172 (Methanol)	Light yellow powdery crystals	-
36	CH ₃	CH ₃		CH ₃		175 - 177 (Tetrahydrofuran)	Light yellow powdery crystals	-
37	CH ₃	CH ₃		CH ₃		197 - 198 (Acetone)	Light yellow powdery crystals	-

(to be continued)

Table 1

(Continued)









Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
38	CH ₃	CH ₃		CH ₃		102.5 - 105 (Methylene chloride)	Light yellow powdery crystals	-
39	CH ₃	CH ₃		C ₂ H ₅		145 - 148 (Methanol)	Light yellow prism-like crystals	-
40	CH ₃	CH ₃				158 - 159 (Chloroform)	Light yellow powdery crystals	-
41	CH ₃	 -CH ₂ OCH ₂ -		CH ₃	CH ₃	121 - 122 (Ether)	Light yellow powdery crystals	-

(to be continued)

0145434

Table 1

(Continued)







Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
42	CH ₃			CH ₃	CH ₃	142 - 145 (Methanol)	Light yellow	1/2·H ₂ O
43	CH ₃			CH ₃	CH ₃	137 - 140 (Methanol)	Yellow powdery crystals	1/2·H ₂ O
44	CH ₃			CH ₃	CH ₃	101.5 - 115 (Ethyl acetate-ether) NMR ¹³	Yellow powdery crystals	HCl·H ₂ O
45	CH ₃	CH ₃		CH ₃		194 - 196 (Methanol)	Light yellow needle-like crystals	-

(to be continued)

0145434

Table 1









(Continued)

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
46	CH ₃	CH ₃		CH ₃		157 - 158 (Methanol)	Light yellow needle-like crystals	-
47	CH ₃	CH ₃		CH ₃		75 - 90 (Ether- n-hexane) NMR ¹⁴⁾	Light yellow powdery crystals	-
48	CH ₃	CH ₃		CH ₃		82 - 85 (Ether- n-hexane)	Light yellow powdery crystals	-

(to be continued)

Table 1









(Continued)

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
49	CH ₃	CH ₃		CH ₃	$\text{-CH}_2\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_2\text{C}\equiv\text{C-}$ 	96 - 102 (Ether- n-hexane)	Light yellow powdery crystals	-
50	CH ₃	CH ₃		CH ₃	$\text{-CH}_2\text{CH}_2\text{-S-CH}_2\text{-}$ 	81.5 - 83 (Ethyl acetate- diisopropyl ether)	Light yellow powdery crystals	-
51	CH ₃	CH ₃		CH ₃	$\text{-CH}_2\text{CH}_2\text{-O-CH}_2\text{-}$ 	77.5 - 78.5 (Ethyl acetate- diisopropyl ether)	Light yellow powdery crystals	-
52	CH ₃	CH ₃		CH ₃	$\text{-CH}_2\text{CH}_2\text{-S-CH}_2\text{-C}\equiv\text{C-}$ 	NMR ¹⁵⁾	-	-

(to be continued)

Table 1







(Continued)

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
53	CH ₃	CH ₃		CH ₃	$-\text{CH}_2\text{CH}_2\text{OCH}_2\text{C}\equiv\text{C}-$ 	NMR ¹⁶⁾	Light yellow indefinite form crystals	-
54	CH ₃	CH ₃		CH ₃	$-\text{CH}_2\text{CH}_2-\text{N}-$ 	115 - 116 (Ethyl acetate-diisopropyl ether)	Light yellow powdery crystals	-
55	CH ₃	CH ₃		CH ₃	$-\text{CH}_2\text{CH}_2-\text{N}-$ 	115 - 116 (Ethyl acetate-diisopropyl ether)	Light yellow powdery crystals	-
56	CH ₃	CH ₃		CH ₃	$-\text{CH}_2\text{CH}_2-\text{N}-$ 	168 - 169 (Acetone)	Light yellow powdery crystals	HCl

(to be continued)

Table 1








(Continued)

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
57	CH ₃	CH ₃		CH ₃		NMR ¹⁷⁾	Yellow indefinite form crystals	-
58	CH ₃	CH ₃		CH ₃		154 - 158 (Chloroform-n-hexane)	Light yellow powdery crystals	2HCl·H ₂ O
59	CH ₃	CH ₃		CH ₃		217 - 220 (Ethyl acetate-methanol-water)	Light yellow needle-like crystals	2HCl

(to be continued)

Table 1

(Continued)

Example No.	R ¹	R ²	R ³	R ⁴	R ⁵	Melting point (°C) (Recrystallization solvent)	Crystal form	Salt
60	CH ₃	CH ₃		CH ₃	$-\text{CH}_2\text{C}\equiv\text{C}-$ 	62 - 68 NMR ¹⁸	Yellow indefinite form crystals	
61	CH ₃	CH ₃		CH ₃	$-\text{CH}_2\text{C}\equiv\text{C}-$ 	144 - 146 (Diisopropyl ether-ethanol)	Light yellow powdery crystals	HCl
62	CH ₃	CH ₃		 CH ₃		NMR ¹⁹	-	-

IR¹⁾ 1660, 1680, 1700 cm⁻¹

NMR¹⁾ (90MHz, CDCl₃):

2.28 (3H, s), 2.35 (6H, s), 3.64 (3H, s), 4.67 (2H, d, J=7Hz), 5.12 (1H, s),
6.06 (1H, broad, s), 6.15 (1H, d, t, Ja=7H, Jb=16Hz), 6.50 (1H, d, J=16Hz),
7.03 (2H, d, J=8Hz), 7.30 (2H, d, J=8Hz), 7.32 (1H, t, J=8Hz), 7.63 (1H, dt,
Ja=2Hz, Jb=8Hz), 7.97 (1H, d, t, Ja=2Hz, Jb=8Hz), 8.12 (1H, t, J=2Hz)

NMR²⁾ (90MHz, CDCl₃):

1.45-2.10 (6H, m), 2.30 (6H, t), 3.40-4.00 (2H, m), 3.56 (3H, s), 4.63 (2H, d,
J=6Hz), 5.10 (1H, s), 5.37 (1H, t, J=3Hz), 6.17 (1H, d, t, Ja=6Hz, Jb=16Hz),
6.45 (1H, d, J=16Hz), 6.65-7.20 (4H, m), 7.30 (1H, t, J=7Hz), 7.62 (1H, d, t,
Ja=2Hz, Jb=7Hz), 7.95 (1H, d, t, Ja=2Hz, Jb=7Hz), 8.13 (1H, t, J=2Hz)

NMR³⁾ (90MHz, CDCl₃):

2.30 (3H, s), 2.32 (3H, s), 3.38 (3H, s), 3.53 (3H, s), 4.65 (2H, d, J=7Hz),
5.10 (1H, s), 5.13 (2H, s), 6.20 (1H, dt, Ja=6Hz, Jb=16Hz), 6.47 (1H, s),
6.87 (1H, d, J=16Hz), 6.87-7.48 (5H, m), 7.62 (1H, dt, Ja=2Hz, Jb=7Hz),
7.93 (1H, dt, Ja=2Hz, Jb=7Hz), 8.12 (1H, t, J=2Hz)

0145434

154

NMR⁴⁾ (90MHz, CDCl₃):

1.50-1.95 (6H, m), 1.97 (3H, s), 2.32 (6H, s), 3.34-3.90 (1H, m), 3.53 (3H, s),
4.65 (2H, d, J=7Hz), 5.03 (1H, s), 5.33 (1H, t, J=3Hz), 5.70 (1H, t, J=7Hz),
5.95 (1H, broad, s), 6.97 (2H, d, J=9Hz), 7.23 (2H, d, J=9Hz), 7.28 (1H, t,
J=7Hz), 7.58 (1H, d, t, Ja=2Hz, Jb=7Hz), 7.93 (1H, dt, Ja=2Hz, Jb=7Hz),
8.08 (1H, t, J=2Hz)

NMR⁵⁾ (90MHz, CDCl₃):

1.30-2.0 (6H, m), 2.32 (6H, s), 3.53 (3H, s), 3.3-3.95 (2H, m), 4.79 (2H, s),
5.07 (1H, s), 5.33 (1H, t, J=3Hz), 6.02 (1H, s), 6.93 (2H, d, J=10Hz), 7.28
(2H, d, J=10Hz), 7.27 (1H, t, J=8Hz), 7.66 (1H, d, J=8Hz), 7.93 (1H, d, J=8Hz),
8.10 (1H, s)

NMR⁶⁾ (90MHz, CDCl₃):

1.40-2.10 (6H, m), 2.32 (6H, s), 3.43 (3H, s), 3.20-3.85 (2H, m), 4.43 (2H, d,
J=7Hz), 4.95 (1H, s), 5.23 (1H, t, J=3Hz), 5.04-5.80 (1H, m), 5.93-6.53 (3H, m),
6.60 (1H, s), 6.87 (2H, d, J=10Hz), 7.20 (2H, d, J=10Hz), 7.23 (1H, t, J=8Hz),
7.53 (1H, d, t, Ja=2Hz, Jb=8Hz), 7.88 (H, d, t, Ja=2Hz, Jb=8Hz)

105

0145434

NMR⁷⁾ (60MHz, CDCl₃):

1.40-2.0 (6H, m), 2.30 (3H, s), 2.33 (3H, s), 3.20-3.90 (2H, m), 3.60 (3H, s),
4.63 (2H, d, J=6Hz), 5.37 (1H, t, J=3Hz), 5.80 (1H, s), 6.03 (1H, d, t, J_a=6Hz,
J_b=16Hz), 6.80 (1H, s), 6.38 (1H, d, J=16Hz), 6.90 (2H, d, J=9Hz), 7.20 (2H,
d, J=9Hz), 6.75-7.80 (4H, m)

NMR⁸⁾ (90MHz, CDCl₃):

1.43-2.15 (6H, m), 2.30 (6H, s), 3.38 (3H, s), 3.20-3.80 (2H, m), 4.47 (2H, dd,
J_a=3Hz, J_b=7Hz), 5.23 (1H, t, J=3Hz), 5.43 (1H, s), 6.83 (1H, broad, s), 5.94
(1H, dt, J_a=6Hz, J_b=16Hz), 6.23 (1H, d, J=16Hz), 6.87 (2H, d, J=9Hz), 7.13 (2H,
d, J=9Hz), 6.75-7.60 (4H, m)

NMR⁹⁾ (90MHz, CDCl₃):

1.5-2.1 (6H, m), 2.28 (3H, s), 3.53 (3H, s), 3.76 (3H, s), 3.4-4.1 (1H, m),
4.57 (2H, d, J=6Hz), 5.27 (1H, t, J=3Hz), 5.99 (1H, dt, J_a=6Hz, J_b=16Hz),
6.07 (1H, s), 6.36 (1H, d, J=16Hz), 6.73 (1H, d, J=7Hz), 6.78 (1H, s), 6.93
(1H, d, J=7Hz), 7.18 (1H, t, J=7Hz), 7.50 (1H, dt, J_a=2Hz, J_b=7Hz), 7.83
(1H, dt, J_a=2Hz, J_b=7Hz), 7.99 (1H, t, J=2Hz)

0145434

NMR¹⁰⁾ (60MHz, CDCl₃):

1.24 (3H, t, J=7Hz), 2.37 (3H, s), 3.66 (3H, s), 3.79 (2H, q, J=7Hz), 4.70 (2H, d, J=6Hz), 5.16 (1H, s), 5.30 (2H, s), 5.91 (1H, broad, s), 6.10 (1H, dt, Ja=6Hz, Jb=16Hz), 6.40 (1H, d, J=16Hz), 7.00-7.47 (4H, m), 7.64 (1H, dt, Ja=2Hz, Jb=7Hz), 7.97 (1H, dt, Ja=2Hz, Jb=7Hz), 8.14 (1H, t, J=2Hz)

NMR¹¹⁾ (90MHz, CDCl₃):

1.40-2.10 (6H, m), 2.26 (6H, m), 3.53 (3H, s), 3.40-4.0 (2H, m), 4.58 (2H, d, J=6Hz), 5.03 (1H, s), 5.30 (1H, t, J=3Hz), 5.97 (1H, dt, Ja=6Hz, Jb=16Hz), 6.30 (1H, broad, s), 6.37 (1H, d, J=16Hz), 6.88 (2H, d, J=9Hz), 7.14 (2H, d, J=9Hz), 7.20 (1H, t, J=6Hz), 7.50 (1H, dt, Ja=2Hz, Jb=6Hz), 7.82 (1H, dt, Ja=2Hz, Jb=6Hz), 8.00 (1H, t, J=2Hz)

NMR¹²⁾ (90MHz, CDCl₃):

1.5-2.1 (6H, m), 2.33 (6H, s), 2.48 (2H, q, J=6Hz), 3.58 (3H, s), 3.5-4.0 (2H, m), 4.15 (2H, t, J=6Hz), 5.09 (1H, s), 5.40 (1H, m), 5.93 (1H, d, t, J=15Hz, 6Hz), 6.30 (1H, s), 6.32 (1H, d, J=15Hz), 6.96 (2H, d, J=9Hz), 7.19 (2H, d, J=9Hz), 7.1-7.3 (1H, m), 7.5-7.7 (1H, d, m, J=6Hz), 7.8-8.0 (1H, D, m, J=6Hz), 8.10 (1H, m)

- NMR¹³) (90MHz, CDCl₃):
- 2.36 (3H, s), 2.40 (3H, s), 3.25 (2H, d, J=6Hz), 3.63 (6H, s), 3.6-3.9 (2H),
5.10 (1H, s), 6.20 (1H, dt, J=16Hz, 6Hz), 6.53 (1H, d, J=16Hz), 7.1-8.3 (10H, m)
- NMR¹⁴) (90MHz, CDCl₃):
- 2.36 (9H, s), 2.75 (2H, t, J=6Hz) 3.53 (2H, s), 3.62 (3H, s), 4.16 (2H, t,
J=6Hz), 5.11 (1H, s), 5.75 (1H, bs), 7.20-8.20 (9H, m)
- NMR¹⁵) (90MHz, CDCl₃):
- 2.33 (6H, s), 2.93 (2H, t, J=6Hz), 3.48 (2H, s), 3.63 (3H, s), 4.30 (2H, t,
J=6Hz), 5.11 (1H, s), 6.22 (1H, s), 7.20-7.50 (6H, m), 7.66 (1H, dt, J=8Hz,
J=2Hz), 7.93 (1H, bd, J=8Hz), 8.12 (1H, t, J=2Hz)
- NMR¹⁶) (90MHz, CDCl₃):
- 2.35 (6H, s), 3.60 (3H, s), 3.76 (2H, t, J=5Hz), 4.17-4.33 (2H, m), 4.36 (2H,
s), 5.11 (1H, s), 5.80 (1H, bs), 7.20-8.20 (9H, m)
- NMR¹⁷) (90MHz, CDCl₃):
- 2.20 (3H, s), 2.27 (3H, s), 2.30 (3H, s), 2.62 (2H, t, J=6Hz), 3.51 (3H, s),
4.18 (2H, m), 5.72 (1H, s), 5.82 (1H, bs), 6.18 (1H, dt, J=6Hz, J=16Hz),
6.43 (1H, d, J=16Hz), 7.10-7.55 (8H, m), 7.62 (1H, d, J=7.5Hz)

NMR¹⁸) (90MHz, CDCl₃):

2.37 (6H, s), 3.64 (3H, s), 4.87 (2H, s), 5.13 (1H, s), 6.20 (1H, brs),
6.94 (1H, dd, J=4.5Hz, J=4Hz), 7.1-7.4 (3H, m), 7.64 (1H, d-m, J=8Hz)
7.97 (1H, d-m J=8Hz), 8.10 (1H, t, J=2Hz)

NMR¹⁹) (90MHz CDCl₃):

1.17 (3H, t, J=7Hz), 1.47 (3H, d, J=5.5Hz), 2.34 (6H, s), 3.35-3.95 (2H, m),
3.60 (3H, s), 4.65 (2H, d= 6Hz), 5.10 (1H, s), 5.35 (1H, 2, J=5.2Hz),
6.05 (1H, dt, J=6Hz, 15.5Hz), 6.46 (1H, d, J=15,5Hz), 6.89 (2H, d, J=9Hz),
7.18-8.09 (6H, m)

1 Example 63

By using suitable starting materials, by a method similar to that described in Example 3, there were prepared compounds of Examples 1 and 7 - 62.

5 Example 64

By using suitable starting materials, by a method similar to that described in Example 5, there were prepared compounds of Examples 1 and 7 - 15, and 17 - 62.

Example 65

- 10 0.5 Gram of methyl 3-(4-hydroxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-pyridine-3,5-dicarboxylate was dissolved in 5 ml of pyridine, then to this solution was added 0.5 ml of acetic anhydride under an ice-cooling condition. The
- 15 reaction mixture was allowed to stand overnight, then poured into an ice-water, and extracted with ether. The ether layer was washed with a diluted hydrochloric acid, 10%-sodium bicarbonate aqueous solution and water in this order, then dried with anhydrous sodium sulfate.
- 20 The ether extract was concentrate, and the residue thus obtained was purified by means of a silica gel column chromatography (eluent: methanol:chloroform=1:100) to yield 0.4 g of methyl 3-(4-acetyloxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-
- 25 pyridine-3,5-dicarboxylate in the form of light yellow indefinite form crystals.

NMR

2.28 (3H, s), 2.35 (6H, s), 3.64 (3H, s),
4.67 (2H, d, J=7Hz), 5.12 (1H, s), 6.06
(1H, broad s), 6.15 (1H, d, t, J_a=7Hz,
J_b=16Hz), 6.50 (1H, d, J=16Hz), 7.03 (2H,
d, J=8Hz), 7.30 (2H, d, J=8Hz), 7.32 (1H,
t, J=8Hz), 7.63 (1H, dt, J_a=2Hz, J_b=8Hz),
7.97 (1H, d, t, J_a=2Hz, J_b=8Hz), 8.12 (1H,
t, J=2H)

1 Example 66

50 Grams of 4-(1-ethoxyethoxy)cinnamyl methyl
2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine was
dissolved in 600 ml of tetrahydrofuran, the solution
5 was cooled to 5°C, then 170 ml of 0.5N-hydrochloric acid
was added in one time to the solution. After 1 hour, the
temperature of the reaction mixture was kept at 10-15°C,
and the reaction mixture was stirred for 2 hours. Sodium
chloride was added to the reaction mixture and the
10 aqueous layer was removed, the 5%-sodium hydrogen carbonate
aqueous solution was added to adjust the pH to 7.0-7.5.
The aqueous layer was removed by separation, then the
organic layer was washed with water, a saturated sodium
chloride aqueous solution in the order, and dried with
15 anhydrous magnesium sulfate. The dried organic layer was
concentrated at 20-25°C, the residue thus obtained was
recrystallized from chloroform to yield 20 g of (4-hydroxy-
phenyl)-2(E)-propenyl methyl 2,6-dimethyl-4-(3-

1 nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate.

Melting point: 145.8 - 149.4°C.

IR: 1670 cm^{-1}

In the above-mentioned Examples 66, the recrystallization was conducted by using ether in place of chloroform, there was yield (4-hydroxyphenyl)-2(E)-propenyl methyl 2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate having the melting point of 163-164°C.
IR: 1660 cm^{-1} , 1680 cm^{-1} .

10 Example 67

3.0 Grams of 3-nitrobenzaldehyde, 2.3 g of methyl 3-aminocrotonate and 5 g of methyl 4(E)-cinnamyloxy-acetoacetate were suspended in 15 ml of isopropanol, and the suspension was refluxed for 10 hours. After the reaction was finished, the solvent was removed by evaporation under a reduced pressure, and the residue thus obtained was purified by means of a silica gel column chromatography (eluent: ether:n-hexane = 1:3 to 1:2). Recrystallization from ether to yield 4.0 g of 3,5-dimethyl 2-cinnamyloxy-6-methyl-4-(3-nitrophenyl)-3,5-dicarboxylate in the form of light yellow powdery substance. Melting point: 121 - 122°C.

By a method similar to that described in Example 67, and by using suitable starting materials, there were prepared compounds of Examples 42-44.

1 Example 68

5.1 Grams of 2-iodomethyl methyl 2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate, and 3.8 g of 4-phenyl-1,2,3,6-tetrahydropyridine were dissolved in 10 ml of toluene, and the solution was refluxed for 6 hours. After the reaction mixture was allowed to stand to cooled at a room temperature, the crystals precipitated in the reaction mixture were removed by filtration, the filtrate was concentrated and the residue thus obtained was dissolved in chloroform. The chloroform layer was washed with a saturated sodium hydrogen carbonate aqueous solution and water in this order, then dried with anhydrous sodium sulfate. Then the dried chloroform extract was concentrated by removing the solvent by evaporation, and the residue thus obtained was purified by means of a silica gel column chromatography (eluent: chloroform:methanol = 300:1), and recrystallized from ethyl acetate-diisopropyl ether to yield 3.6 g of methyl 2-(4-phenyl-1,2,3,6-tetrahydropyridyl)ethyl 2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate in the form of light yellow powdery substance. Melting point: 115 - 116°C.

By a method similar to that described in Example 68, and by using suitable starting materials, there were prepared compounds of Examples 47 - 59.

Example 69

10.2 Grams of methyl 2-piperazinylethyl 2,6-

1 dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-
dicarboxylate and 4.5 g of 3-phenylpropargyl bromide in
100 ml of acetonitrile were stirred at 30°C for 5 hours
under heating condition. After the reaction was
5 finished, chloroform was added to the reaction mixture,
and the mixture was washed with a saturated sodium
hydrogen carbonate aqueous solution and water in this
order, and dried with anhydrous sodium sulfate. The
dried chloroform extract was concentrated by removing
10 the solvent by evaporation, the residue thus obtained
was purified by means of a silica gel column chromato-
graphy (eluent: dichloromethane - methanol:dichloromethane
= 1:100). The product thus purified was then converted
into a hydrochloride by adding hydrochloric acid-dioxane,
15 and recrystallized from ethyl acetate-methanol-water to
yield 1.2 g of methyl 2-[4-(3-phenyl-2-propynyl)-1-
piperazinyl]ethyl 2,6-dimethyl-4-(3-nitrophenyl)-1,4-
dihydropyridine-3,5-dicarboxylate in the form of light
yellow needle-like crystals. Melting point: 217 - 220°C.

20 By a method similar to that described in Example
69, and by using a suitable starting material, there was
prepared a compound of Example 58.

Pharmacological tests

The results of the pharmacological test on
25 dihydropyridine derivatives of the present invention are
shown below.

The test compounds used in the tests are as

1 follows.

Test
Compound
No.

1. Methyl 3-(4-hydroxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
2. Methyl 3-phenyl-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
3. Methyl 5-(4-hydroxyphenyl)-2(E),4(E)-pentadienyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
4. Methyl 3-methyl-3-(4-hydroxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
5. Methyl 3-phenyl-2-propynyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
6. Methyl 3-phenylpropyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate [Reference compound: Japanese Patent Application Kokai (Laid-open) No. 56-36455 (1981)]
7. Methyl 3-(3-methoxy-4-hydroxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
8. Methyl 3-(4-methoxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dihydrocarboxylate
9. Methyl 3-(1-methyl-1,2,3,4-tetrazol-5-yl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
10. Methyl 3-furyl-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
11. Methyl 3-(3-chloro-4-hydroxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
12. Methyl 3-(4-hydroxyphenyl)-2-propynyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate

13. Methyl 3-(2-hydroxyphenyl)-2(E)-propenyl
1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-
pyridine-3,5-dicarboxylate
14. Methyl 3-(4-hydroxyphenyl)-2(E)-propenyl
1,4-dihydro-2,6-dimethyl-4-(2-trifluoro-
methylphenyl)pyridine-3,5-dicarboxylate
15. Methyl 4-(4-methylthiophenyl)-3(E)-butenyl
1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-
pyridine-3,5-dicarboxylate
16. Methyl 3-(4-acetyloxyphenyl)-2(E)-propenyl
1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-
pyridine-3,5-dicarboxylate
17. Methyl 2-[4-(4-methylphenyl)-1,2,3,6-tetra-
hydropyridyl]ethyl 1,4-dihydro-2,6-dimethyl-
4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
18. Methyl 3-(4-hydroxyphenyl)-2(E)-propenyl
1,4-dihydro-2,6-dimethyl-4-phenylpyridine-
3,5-dicarboxylate
19. Methyl 3-(4-hydroxyphenyl)-2(E)-propenyl
1,4-dihydro-2,6-dimethyl-4-(2-methylphenyl)-
pyridine-3,5-dicarboxylate
20. Methyl 3-(4-hydroxyphenyl)-2(E)-propenyl
1,4-dihydro-2,6-dimethyl-4-(3,4-dimethoxy-
phenyl)pyridine-3,5-dicarboxylate
21. Dimethyl 1,4-dihydro-2-[3-(4-hydroxyphenyl)-
2(E)-propenyloxymethyl]-6-methyl-4-(3-
nitrophenyl)pyridine-3,5-dicarboxylate
22. Methyl 3-(2-thienyl)-2-propynyl 1,4-dihydro-
2,6-dimethyl-4-(3-nitrophenyl)pyridine-
3,5-dicarboxylate
23. Methyl 2-[4-(3-phenyl-2-propynyl)-1-
piperazinyl]ethyl 1,4-dihydro-2,6-dimethyl-
4-(3-nitrophenyl)pyridine-3,5-dicarboxylate
24. Methyl 3-(4-hydroxyphenyl)-2(E)propenyl
1,4-dihydro-2,6-dimethyl-4-(2,4-dichloro-
phenyl)pyridine-3,5-dicarboxylate
25. Methyl 3-(3-hydroxyphenyl)-2(E)propenyl
1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-
pyridine-3,5-dicarboxylate
26. Methyl 2-(N-methyl-N-benzylamino)ethyl 1,4-
dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-
dicarboxylate [Reference compound known as
"Nicardipine"]

1 Pharmacological test-1

Vasodilative effects of the dihydropyridine derivatives of the present invention were determined by measuring the systolic blood pressures of test
5 animals before and after the administration of test compound.

Systolic blood pressures and heart beat of SHR-rats (spontaneously hypertensive rats) were determined by a "tail cuff method". Thus the test animal was
10 placed in a thermostat chamber (Type: FR-12RS, manufactured by Isuze & Co.) and was warmed at 40°C for 15 minutes so as to vasodilate the tail arteria, the systolic blood pressures were measured by an electro sphygmomanometer (Type: PE-300, manufactured by Narco-
15 Biosystems, Inc.) and recorded by an ink-writing recorder (Type: RECTI-HORIZ 8s, manufactured by San-Ei Instrument & Co.). The experiments were conducted under non-anesthetized and semi-confinement conditions. The test compound was orally administered forcedly by
20 using a sonde for oral administration. The test compound was suspended in 0.15%-gum arabic powder aqueous solution so as to make the quantity of the test compound to 2.5 ml/kg. The test animal was not fasted, and the systolic blood pressures (mm-Hg) were measured before the
25 administration (hereinafter referred to as "PRE") and after the administration (8, 24, 30, 48, 54 and 72 hours after the administration) of the test compound. The data of systolic blood pressure measured before the

1 administration are shown in absolute value of mmHg, and the data measured after the administration are shown in the differences from the absolute values. The results are shown in Table 2 as follows.

Table 2

Test Compound No.	Dosage (mg/kg) p.o.	PRE	8	Blood pressure (mm-Hg) After the administration (Hours)				
				24	30	48	54	72
1	30	195.0	-	-41.0	-52.0	-26.5	-19.8	-12.5
2	30	187.4	-	-38.6	-21.2	-21.0	-19.2	-
3	30	180.8	-	-31.0	-30.8	-15.6	-12.6	-
4	30	184.4	-	-18.0	-25.8	-19.8	-14.6	-
5	30	180.8	-	-75.4	-66.8	-20.8	-10.6	-9.6
6	30	185.0	-20.3	+1.8	-	-	-	-

1 As can be seen the data shown in Table 2, the
vasodilative effects of the dihydropyridine derivatives
of the present invention can be prolonged for certain
length of period as compared with that of the reference
5 compounds.

Pharmacological tests

Calmodulin (calcium-dependent modulator protein)
inhibitory activity of each of the test compounds was
determined by judging from the difference between IC_{50}
10 (50%-inhibitory concentration) of calmodulin inhibitory
activity of the test compound being measured in the
presence of calmodulin together with calmodulin-dependent
cyclic-AMP phosphodiesterase, and another IC_{50} of
calmodulin-dependent cyclic-AMP phosphodiesterase inhibi-
15 tory activity of the same test compound being measured
in the absence of calmodulin. In other words, calmodulin-
specific-inhibitory activity of each of the test compounds
was determined in case that the test compound shows
higher inhibitory activity against calmodulin-dependent
20 cyclic-AMP phosphodiesterase in the presence of calmodulin
than another inhibitory activity against calmodulin-
dependent cyclic-AMP phosphodiesterase only.

1) Reagents used in the test

(1) Calmodulin: A calmodulin product manu-
25 factured by Amano & Co., which was isolated from the
brain of bovine, and purified so as to be considered as
a single substance with respect to SDS-page (poly-

1 acrylamide gel electrophoresis) method.

(2) Calmodulin-dependent cyclic-AMP phosphodiesterase (EC 3.1.4.17): An enzyme substance isolated from the heart of bovine and purified partially by a method of modified version of the disclosure in H. C. Ho, T. S. Teo, et al.: "Biochim. Biophys. Acta", 429, 461 (1976).

(3) 5'-Nucleotidase (EC 3.1.3.5): Grade IV substance (isolated from Crotalus adamanteus venom) manufactured by Sigma & Co.

(4) Others: Remainder of the reagents used in the test were those of reagent grade chemicals manufactured by Wako Pure-Chemical Industries, Ltd.

2) Method for the test

15 Calmodulin inhibitory activity of each of the test compounds was measured by a method of modified version of the disclosure in T. S. Teo and T. H. Wang: "J. Bio. Chem.", 248, 588 (1973).

(1) Cyclic-AMP-phosphodiesterase: One unit thereof hydrolyzes 1.0 micromole of 3':5'-cyclic-AMP to 5'-AMP per minute at pH 7.5 at 30°C, in the presence of a saturating level of calmodulin.

(2) Calmodulin: One unit thereof stimulates 0.015 activated unit of cyclic-AMP phosphodiesterase to 25 50% of the maximum activity of the enzyme.

(3) 5'-Nucleotidase: One unit thereof hydrolyzes 1.0 micromole of inorganic phosphorus from adenosine 5'-monophosphate per minute at pH 9.0 at 37°C.

1 3) Reactions in the tests

(1) Calmodulin-cyclic-AMP phosphodiesterase
inhibitory activity:

40 mM of Imidazol, 20 mM of $MgCl_2$,
5 20 mM of $CaCl_2$, 0.008 unit of cyclic-AMP phospho-
diesterase, 1.0 unit of calmodulin, 0.2 unit of 5'-
nucleotidase and 1.0 ml of 10 mM-tris(hydroxymethyl)amino-
methane/HCl buffer solution (pH 7.0) containing 0.5 mM
of cyclic-AMP were mixed together, and reacted at 30°C
10 for 30 minutes. Each of the test compounds was dis-
solved in methanol or N,N-dimethylformamide as the
solvent, provided that the quantity of the solvent was
not exceed 2% of the total amount of the mixture. After
the reaction was completed, the reaction mixture was
15 ice-cooled, and 0.5 ml each of aqueous solutions of
16.5%-trichloroacetic acid, 1%-thiourea, 3%-ammonium
ferrous sulfate were respectively added to the reaction
mixture. Further, 0.15 ml of 4.4%-ammonium molybdate
solution was added to the mixture and the whole of the
20 mixture was stirred, and was centrifuged at 3,000 r.p.m.
for 10 minutes. Then, the centrifuged mixture was
allowed to stand at a room temperature for 20 minutes.
The $OD_{660\text{ nm}}$ (optical density at 660 nm) was measured.

(2) Cyclic-AMP phosphodiesterase inhibitory
25 activity:

The reaction was conducted by a method
similar to that described in (1) as mentioned above,
except that 1 mM of EGTA [ethylene glycol-bis(β -amino

1 ethyl ether)-N,N-tetraacetic acid] was used in place of 20 mM of CaCl_2 . The reaction was conducted for 3 hours. The results are shown in Table 3 as follows.

Table 3

<u>Test Compound No.</u>	<u>Calmodulin-cyclic-AMP phosphodiesterase IC₅₀ (μg/ml)</u>	<u>Cyclic-AMP phosphodiesterase IC₅₀ (μg/ml)</u>
1	4.1	60
2	2.5	>100
3	0.85	>100
7	5.2	86
8	2.7	>100
9	5.3	36
10	1.55	>100
11	5.3	100
12	0.78	13
13	1.65	12.5
14	6.25	>100
15	0.062	0.23
16	3.0	50
17	5.5	>100
18	6.8	32
19	5.8	58
20	3.2	30
21	6	>100
22	1.1	>100
23	8.0	>100
24	2.45	15

0145434


- 124 -

25	0.9	24.6
26	6.25	13

1 As can be seen from the data shown in Table 3.
the dihydropyridine derivatives of the present invention
have specific inhibitory activity against calmodulin
as compared with that of indicated by known compound.

WHAT IS CLAIMED IS:

1. A dihydropyridine derivative represented by the general formula,



wherein R^1 and R^4 are each a lower alkyl group; R^2 is a lower alkyl group or a group of the formula $-\text{CH}_2-\text{A}-\text{R}^6$ [wherein A is a straight-chain or branched-chain unsaturated hydrocarbon residual group which may have an oxygen atom or a group of the formula $-\text{N}-\text{R}^7$ (wherein R^7 is a lower alkyl group)]; and R^6 is a phenyl group which may have a hydroxyl group]; R^3 is a phenyl group which may have 1 to 2 substituents selected from the group consisting of a nitro group, a lower alkyl group which may have 1 to 3 halogen atoms, a lower alkoxy group and a halogen atom; and R^5 is a lower alkyl group, a 1,2,3,6-tetrahydropyridyl-lower alkyl group which may have, as the substituent, a phenyl group which may have halogen atoms or lower alkyl groups as the substituents on the phenyl ring, or a group of the formula $-\text{CH}_2-\text{A}'-\text{R}^8$ [wherein A' is a straight-chain or branched-chain unsaturated hydrocarbon residual group which may have or may not have an oxygen atom, a sulfur atom, a group of the formula $-\text{N}-\text{R}^7$ (wherein R^7 is a lower alkyl group)],

or a group of the formula $\text{-N} \begin{array}{|c|} \hline \text{ } \\ \hline \end{array} \text{N-}$ in the unsaturated hydrocarbon residual group; and R^8 is a phenyl group which may have 1 to 3 substituents selected from the group consisting of a lower alkoxy group, a halogen atom, a lower alkylthio group, a hydroxyl group; a lower alkanoyloxy group, a tetrahydropyranyloxy group and a lower alkoxy-lower alkoxy group; a pyridyl group, a thienyl group, a furyl group, or a tetrazolyl group which may have a lower alkyl group as the substituent] provided that when R^5 is a lower alkyl group, then R^2 should be of a group of the formula $\text{-CH}_2\text{-A-R}^6$ (wherein A and R^6 are the same as defined above) or a salt thereof.

2. The dihydropyridine derivatives according to Claim 1, wherein R^2 is a group of the formula $\text{-CH}_2\text{-A-R}^6$ [wherein A is a straight-chain or branched-chain unsaturated hydrocarbon residual group having 2 to 6 carbon atoms in the unsaturated hydrocarbon residual moiety, having 1 to 3 double bonds and/or triple bonds therein, said unsaturated hydrocarbon residual group may have or may not have an oxygen atom or a group of the formula -N-R^7 (wherein R^7 is a lower alkyl group); and R^6 is a phenyl group which may have a hydroxy group as the substituent].

3. The dihydropyridine derivatives according to Claim 1, wherein R^2 is an alkyl group having 1 to 6 carbon atoms.

4. The dihydropyridine derivatives according to

Claim 2, wherein R^5 is an alkyl group having 1 to 6 carbon atoms.

5. The dihydropyridine derivatives according to Claim 3, wherein R^5 is a 1,2,3,6-tetrahydropyridyl- C_{1-6} -alkyl group which may have, as the substituent on the pyridyl ring, a phenyl group which may have halogen atoms or lower alkyl groups as the substituents on the phenyl ring.

6. The dihydropyridine derivatives according to Claim 3, wherein R^5 is a group of the formula $-CH_2-A'-R^8$ [wherein A' is a straight-chain or branched-chain unsaturated hydrocarbon residual group having 2 to 6 carbon atoms in the unsaturated hydrocarbon residual moiety, having 1 to 3 double bonds and/or triple bonds therein, said unsaturated hydrocarbon residual group may have or may not have an oxygen atom, a sulfur atom, a group of the formula $-N-R^7$ (wherein R^7 is a C_{1-6} -alkyl group)], or a group of the formula $-N \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} N-$ in the unsaturated hydrocarbon residual group.

7. The dihydropyridine derivatives according to Claim 2, wherein R^5 is a 1,2,3,6-tetrahydropyridyl- C_{1-6} -alkyl group which may have, as the substituent on the pyridyl ring, a phenyl group which may have halogen atoms or C_{1-6} -alkyl groups as the substituents on the phenyl ring; or a group of the formula $-CH_2-A'-R^8$ (wherein A' and R^8 are the same as defined above)

8. The dihydropyridine derivatives according to Claim 6, wherein A' is a straight-chain or branched-chain

unsaturated hydrocarbon residual group having 2 to 6 carbon atoms in the unsaturated hydrocarbon residual moiety, which contains an oxygen atom, a sulfur atom, or a group of the formula -N-R^7 (wherein R^7 is C_{1-6} -alkyl group), or a group of the formula $\text{-N} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{N-}$ in the unsaturated hydrocarbon residual group.

9. The dihydropyridine derivatives according to Claim 6, wherein A' is a straight-chain or branched-chain unsaturated hydrocarbon residual group having 2 to 6 carbon atoms in the unsaturated hydrocarbon residual moiety, which does not contain an oxygen atom, a sulfur atom, or a group of the formula -N-R^7 (wherein R^7 is C_{1-6} -alkyl group), or a group of the formula $\text{-N} \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \text{N-}$ in the unsaturated hydrocarbon residual group.

10. The dihydropyridine derivatives according to Claim 8, wherein R^8 is a phenyl group which may have 1 to 3 substituents on the phenyl ring, selected from the group consisting of 2 C_{1-6} -alkoxyl group, a halogen atom, a C_{1-6} -alkylthio group, a hydroxyl group, C_{1-6} -alkanoyloxy group, a tetrahydropyranyloxy group and C_{1-6} -alkoxy- C_{1-6} -alkoxy group.

11. The dihydropyridine derivatives according to Claim 9, wherein R^8 is a phenyl group which may have 1 to 3 substituents on the phenyl ring, selected from the group consisting of a C_{1-6} -alkoxy group, a halogen atom, a C_{1-6} -alkylthio group, a hydroxyl group, a C_{1-6} -alkanoyloxy group, a tetrahydropyranyloxy group and a C_{1-6} -alkoxy- C_{1-6} -alkoxy group.

12. The dihydropyridine derivatives according to Claim 8 or 9, wherein R^8 is a pyridyl group, a thienyl group, a furyl group, or a tetrazolyl group which may have a lower alkyl group as the substituent.
13. The dihydropyridine derivatives according to Claim 11, wherein R^8 is a phenyl group which may have 1 to 3 substituents on the phenyl ring, selected from the group consisting of a hydroxyl group and C_{1-6} -alkanoyloxy group.
14. The dihydropyridine derivatives according to Claim 11, wherein R^8 is a phenyl group which contains 1 to 3 substituents on the phenyl ring, selected from the group consisting of a C_{1-6} -alkoxy group, a halogen atom, a C_{1-6} -alkylthio group, a tetrahydropyranyloxy group and C_{1-6} -alkoxy- C_{1-6} -alkoxy group.
15. The dihydropyridine derivatives according to Claim 13 or 14, wherein R^3 is a phenyl group which may have 1 to 2 nitro groups as the substituents on the phenyl ring.
16. The dihydropyridine derivatives according to Claim 13 or 14, wherein R^3 is a phenyl group having 1 to 2 C_{1-6} -alkyl group as the substituents on the phenyl ring, said C_{1-6} -alkyl group contains 1 to 3 halogen atoms as the substituents.
17. The dihydropyridine derivatives according to Claims 13 or 14, wherein R^3 is a phenyl group having 1 to 2 substituents on the phenyl ring, selected from the group consisting of a C_{1-6} -alkyl group, a C_{1-6} -alkoxy

group and a halogen atom.

18. The dihydropyridine derivatives according to Claim 15, wherein the straight-chain or branched-chain unsaturated hydrocarbon residual group is a straight-chain or branched-chain unsaturated hydrocarbon residual group which only contains 1 - 3 double bonds.


19. The dihydropyridine derivatives according to Claim 15, wherein the straight-chain or branched-chain unsaturated hydrocarbon residual group is a straight-chain or branched-chain unsaturated hydrocarbon residual group which only contains 1 - 3 triple bonds.

20. Methyl 3-(4-hydroxyphenyl)-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate.

21. Methyl 3-phenyl-2(E)-propenyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)pyridine-3,5-dicarboxylate.


22. Methyl 3-phenyl-2-propynyl 1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-pyridine-3,5-dicarboxylate.

23. A process for preparing a dihydropyridine derivative represented by the general formula (1),



[wherein R^1 , R^2 , R^3 , R^4 and R^5 are the same as defined above], by reacting a compound represented by the

general formula (2),




[wherein R^1 , R^2 , R^3 and R^4 are the same as defined above],
with a compound represented by the general formula (3),



[wherein X is a hydroxyl group or a halogen atom].

24. A process for preparing a dihydropyridine derivative represented by the general formula (1),



[wherein R^1 , R^2 , R^3 , R^4 and R^5 are the same as defined above], by reacting a compound represented by the general formula (4),




[wherein R^2 and R^5 are the same as defined above], with

0145434


a compound represented by the general formula (5),



[wherein R^3 is the same as defined above] to prepare a compound represented by the general formula




[wherein R^2 , R^3 and R^5 are the same as defined above], then thus prepared compound represented by the general formula (6) is reacted with a compound represented by the general formula (7),



[wherein R^1 and R^4 are the same as defined above] to prepare the desired dihydropyridine derivative represented by the general formula (1).


25. A process for preparing a dihydropyridine derivative represented by the general formula (1a),

0145434



(1a)

[wherein R^1 , R^2 , R^3 and R^4 are the same as defined above; E is a lower alkylene group; R^{14} is a 1,2,3,6-tetrahydropyridyl group which may have, as the substituent, a phenyl group which may have halogen atoms or lower alkyl groups as the substituents on the phenyl ring; or a group of the formula $R^6-(D')_0-Y-$ (wherein R^6 is the same as defined above; D' is a saturated- or unsaturated-alkylene group; 0 is 0 or 1; Y is an oxygen atom, a sulfur atom, a group of the formula $-N-R^7$ (wherein R^7 is a lower alkyl group), or a group of the formula $-N \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} N-$], reacting a compound represented by the general formula (31),




(31)


[wherein R^1 , R^2 , R^3 , R^4 and E are the same as defined above; and X^1 is a halogen atom] with a compound represented by the general formula (32),

[wherein R¹⁴ is the same as defined above] to prepare the desired dihydropyridine derivative represented by the general formula (1a).

26. A process for preparing a dihydropyridine derivative represented by the general formula (1b),



[wherein R¹, R², R³, R⁴, D', _O E and R⁸ are the same as defined above], by reacting a compound represented by the general formula (36),



[wherein R¹, R², R³, R⁴ and E are the same as defined above], with a compound represented by the general formula (34a),



[wherein R^8 is the same as defined above; and X^2 is a halogen atom] to prepare the desire dihydropyridine derivative represented by the general formula (1b).

27. A hypotensive composition containing, as the active ingredient a dihydropyridine derivative represented by the general formula (1) as claimed in Claim 1 and pharmaceutically acceptable carriers.

28. A coronary blood flow improving composition containing, as the active ingredient a dihydropyridine derivative represented by the general formula (1) as claimed in Claim 1 and pharmaceutically acceptable carriers.